

ISSN 2518-1467 (Online),
ISSN 1991-3494 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Х А Б А Р Ш Ы С Ы

ВЕСТНИК

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

THE BULLETIN

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

1944 ЖЫЛДАН ШЫҒА БАСТАҒАН
ИЗДАЕТСЯ С 1944 ГОДА
PUBLISHED SINCE 1944

2

АЛМАТЫ
АЛМАТЫ
ALMATY

2018

MARCH
МАРТ
HAУРЫЗ

NAS RK is pleased to announce that Bulletin of NAS RK scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of Bulletin of NAS RK in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential multidiscipline content to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабаршысы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабаршысының Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді мультидисциплинарлы контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Вестник НАН РК» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Вестника НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному мультидисциплинарному контенту для нашего сообщества.

Б а с р е д а к т о р ы

х. ғ. д., проф., ҚР ҰҒА академигі

М. Ж. Жұрынов

Р е д а к ц и я а л қ а с ы:

Абиев Р.Ш. проф. (Ресей)
Абишев М.Е. проф., корр.-мүшесі (Қазақстан)
Аврамов К.В. проф. (Украина)
Аппель Юрген проф. (Германия)
Баймуқанов Д.А. проф., корр.-мүшесі (Қазақстан)
Байпақов К.М. проф., академик (Қазақстан)
Байтулин И.О. проф., академик (Қазақстан)
Банас Иозеф проф. (Польша)
Берсимбаев Р.И. проф., академик (Қазақстан)
Велихов Е.П. проф., РҒА академигі (Ресей)
Гашимзаде Ф. проф., академик (Әзірбайжан)
Гончарук В.В. проф., академик (Украина)
Давлетов А.Е. проф., корр.-мүшесі (Қазақстан)
Джрбашян Р.Т. проф., академик (Армения)
Қалимолдаев М.Н. проф., академик (Қазақстан), бас ред. орынбасары
Лаверов Н.П. проф., академик РАН (Россия)
Лупашку Ф. проф., корр.-мүшесі (Молдова)
Мохд Хасан Селамат проф. (Малайзия)
Мырхалықов Ж.У. проф., академик (Қазақстан)
Новак Изабелла проф. (Польша)
Огарь Н.П. проф., корр.-мүшесі (Қазақстан)
Полещук О.Х. проф. (Ресей)
Поняев А.И. проф. (Ресей)
Сагиян А.С. проф., академик (Армения)
Сатубалдин С.С. проф., академик (Қазақстан)
Таткеева Г.Г. проф., корр.-мүшесі (Қазақстан)
Умбетаев И. проф., академик (Қазақстан)
Хрипунов Г.С. проф. (Украина)
Юлдашбаев Ю.А. проф., РҒА корр.-мүшесі (Ресей)
Якубова М.М. проф., академик (Тәжікстан)

«Қазақстан Республикасы Ұлттық ғылым академиясының Хабаршысы».

ISSN 2518-1467 (Online),

ISSN 1991-3494 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы»РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5551-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 2000 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
www: nauka-nanrk.kz, bulletin-science.kz

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2018

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д. х. н., проф. академик НАН РК
М. Ж. Журинов

Р е д а к ц и о н н а я к о л л е г и я:

Абиев Р.Ш. проф. (Россия)
Абишев М.Е. проф., член-корр. (Казахстан)
Аврамов К.В. проф. (Украина)
Апель Юрген проф. (Германия)
Баймуканов Д.А. проф., чл.-корр. (Казахстан)
Байпаков К.М. проф., академик (Казахстан)
Байтулин И.О. проф., академик (Казахстан)
Банас Иозеф проф. (Польша)
Берсимбаев Р.И. проф., академик (Казахстан)
Велихов Е.П. проф., академик РАН (Россия)
Гашимзаде Ф. проф., академик (Азербайджан)
Гончарук В.В. проф., академик (Украина)
Давлетов А.Е. проф., чл.-корр. (Казахстан)
Джрбашян Р.Т. проф., академик (Армения)
Калимолдаев М.Н. академик (Казахстан), зам. гл. ред.
Лаверов Н.П. проф., академик РАН (Россия)
Лупашку Ф. проф., чл.-корр. (Молдова)
Мохд Хасан Селамат проф. (Малайзия)
Мырхалыков Ж.У. проф., академик (Казахстан)
Новак Изабелла проф. (Польша)
Огарь Н.П. проф., чл.-корр. (Казахстан)
Полещук О.Х. проф. (Россия)
Поняев А.И. проф. (Россия)
Сагиян А.С. проф., академик (Армения)
Сатубалдин С.С. проф., академик (Казахстан)
Таткеева Г.Г. проф., чл.-корр. (Казахстан)
Умбетаев И. проф., академик (Казахстан)
Хрипунов Г.С. проф. (Украина)
Юлдашбаев Ю.А. проф., член-корр. РАН (Россия)
Якубова М.М. проф., академик (Таджикистан)

«Вестник Национальной академии наук Республики Казахстан».

ISSN 2518-1467 (Online),

ISSN 1991-3494 (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5551-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год

Тираж: 2000 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18.

www: nauka-nanrk.kz, bulletin-science.kz

© Национальная академия наук Республики Казахстан, 2018

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

E d i t o r i n c h i e f

doctor of chemistry, professor, academician of NAS RK

M. Zh. Zhurinov

E d i t o r i a l b o a r d:

Abiyev R.Sh. prof. (Russia)
Abishev M.Ye. prof., corr. member. (Kazakhstan)
Avramov K.V. prof. (Ukraine)
Appel Jurgen, prof. (Germany)
Baimukanov D.A. prof., corr. member. (Kazakhstan)
Baipakov K.M. prof., academician (Kazakhstan)
Baitullin I.O. prof., academician (Kazakhstan)
Joseph Banas, prof. (Poland)
Bersimbayev R.I. prof., academician (Kazakhstan)
Velikhov Ye.P. prof., academician of RAS (Russia)
Gashimzade F. prof., academician (Azerbaijan)
Goncharuk V.V. prof., academician (Ukraine)
Davletov A.Ye. prof., corr. member. (Kazakhstan)
Dzhrbashian R.T. prof., academician (Armenia)
Kalimoldayev M.N. prof., academician (Kazakhstan), deputy editor in chief
Laverov N.P. prof., academician of RAS (Russia)
Lupashku F. prof., corr. member. (Moldova)
Mohd Hassan Selamat, prof. (Malaysia)
Myrkhalykov Zh.U. prof., academician (Kazakhstan)
Nowak Isabella, prof. (Poland)
Ogar N.P. prof., corr. member. (Kazakhstan)
Poleshchuk O.Kh. prof. (Russia)
Ponyaev A.I. prof. (Russia)
Sagiyani A.S. prof., academician (Armenia)
Satubaldin S.S. prof., academician (Kazakhstan)
Tatkeyeva G.G. prof., corr. member. (Kazakhstan)
Umbetayev I. prof., academician (Kazakhstan)
Khripunov G.S. prof. (Ukraine)
Yuldashbayev Y.A., prof. corresponding member of RAS (Russia)
Yakubova M.M. prof., academician (Tadjikistan)

Bulletin of the National Academy of Sciences of the Republic of Kazakhstan.

ISSN 2518-1467 (Online),

ISSN 1991-3494 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5551-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 2000 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/>, <http://bulletin-science.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2018

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

**V. G. Semenov¹, D. A. Baimukanov², N. I. Kosyaev¹, R. M. Mudarisov³,
N. I. Morozova⁴, F. A. Musayev⁴, D. A. Nikitin¹, M. B. Kalmagambetov²**

¹Chuvash state agricultural academy, Cheboksary, Chuvash Republic, Russia,

²Kazakh research institute of livestock production and forage production, Almaty, Kazakhstan,

³Bashkir state agricultural university, Ufa, Republic of Bashkortostan, Russia,

⁴Ryazan state agrotechnological university named after P. A. Kostychev, Ryazan, Ryazan region, Russia.

E-mail: semenov_v.g@list.ru dbaimukanov@mail.ru kocyevni81@mail.ru r-mudarisov@mail.ru

morozova@rgatu.ru musayev@rgatu.ru nikitin_d_a@mail.ru animal_feeding@mail.ru

GROWTH, DEVELOPMENT AND MEAT QUALITIES OF BULL-CALVES AGAINST THE BACKGROUND OF APPLICATIONS WITH BIOLOGICAL PREPARATIONS OF THE PREVENTION SERIES

Abstract. For the first time on the basis of complex studies the zootechnical expediency of the developed biological preparations Prevention-N-A and Prevention-N-E in beef production technology for the realization of bioresource potential of meat qualities of Black Motley bull-calves was scientifically and experimentally proved. Activation of the growth and development of bull-calves in the periods of growth, rearing and fattening was established against the background of biopreparations, which resulted in higher slaughter and meat qualities of the carcasses and, as a consequence, in a higher yield of valuable cuts: brisket and sirloin by 6.1 and 4.0 kg ($P < 0,01-0,001$), rump by 2,6 and 1,7 kg ($P < 0,05-0,01$) and thick flank by 8,6 and 7,1 kg ($P < 0,001$) (compared with the control group).

The largest content of highest-quality meat came from the carcasses of the bull calves of the 1st ($27,8 \pm 0,72$ kg) and the 2nd ($26,7 \pm 0,58$ kg) test groups: 3.5 and 2.4 kg more, respectively, as compared with the control group ($24,3 \pm 0,73$ kg), and also from their cuts: brisket and sirloin - 0,9 and 0,7 kg more, respectively, rump - 0,5 and 0,3 kg, and thick flank - 2,3 and 1, 5 kg ($P < 0.05-0.001$).

The high quality of meat carcasses by organoleptic, biochemical and spectrometric indicators and, consequently, the safety of the tested preparations was proved. It was found that biological preparations lead to the realization of bioresource potential of the organism due to activation of haemopoiesis, cellular and humoral factors of non-specific resistance (with a more pronounced Prevention-N-A effect). The novelty of the data obtained is confirmed by the patents of the Russian Federation for invention No. 2602687 and No. 2622765 registered in the Public Register of Inventions of the Russian Federation on October 26, 2016, and June 19, 2017, respectively.

Keywords: bull-calves, growth, rearing, fattening, biopreparations Prevention-N-A and Prevention-N-E, meat qualities.

Introduction. Russian cattle breeding is one of the main life-supporting sectors of domestic agrarian production, which has a decisive influence on the level of the country's food supply and determines the health of the nation.

The industry development strategy is aimed at increasing the share of domestic production, forming dairy and meat resources in accordance with scientifically based consumption standards, increasing its competitiveness and investment attractiveness, and finding solution to the most important socio-economic task of providing the population with biologically wholesome products [2, 4, 11, 17, 21].

In terms of production volumes, the domestic cattle-breeding industry lags behind the targets by 25%, with more than 95% of beef produced by slaughtering replacements and the rejected adult livestock of dairy and combined productivity, the slaughter contingent of which and the level of productivity do not

provide the necessary production volumes [1, 5, 9, 15, 20, 29, 30]. In most regions of Russia, Chuvashia included, the predominant dairy cattle breed remains Black Motley (55.7%), as the most highly productive with good feed payment. As a result of the selection, the cattle acquired features inherent in the dairy type, but with good meat qualities, and therefore has a great potential for productivity, overtopping many other breeds by zootechnical and economic indicators. Thus, for beef production young Black Motley calves are most commonly used, being more adapted and maximally realizing the bioresource potential under optimal feeding and maintenance conditions [6, 22, 24].

In order to prevent immunodeficiency, to stimulate the level of non-specific resistance of the organism a wide range of feed and bioactive additives, immunocorrectors, antioxidants and biopreparations are used to fight environmental and technological stress factors and to realize the bioresource potential of the meat qualities of bull-calves; however, many of them do not exhibit the desired bioeffect [3, 7, 13, 14, 16, 18, 19, 26, 34, 35].

In this context, the development and introduction of complex biologics into the beef production technology to activate the protective-adaptive functions of the organism to the habitat conditions and to realize the bioresource potential of the meat qualities of bull-calves is an actual problem of modern zootechnical science and practice [19].

The aim of the present study is to realize the bioresource potential of meat qualities of Black Motley bull-calves by using biopreparations Prevention-N-A and Prevention-N-E.

Materials and methods. The experimental research was carried out on the dairy farm (an agricultural production cooperative) 'Novyi Put' of the Alikovsky District of the Chuvash Republic in accordance with the plan of scientific research of Chuvash State Agricultural Academy, and the processing of materials was carried out in the Chuvash Republican Veterinary Laboratory of the State Veterinary Service of the Chuvash Republic; in the laboratory of bio- and nanotechnologies and in the laboratory of the Department of morphology, obstetrics and therapy of Chuvash State Agricultural Academy in the period from 2013 to 2017.

The objects of the research were three groups of Black Motley bull-calves (15 animals each), from birth to 540 days of age. Newborn bull-calves of all the groups were kept with their mother in the maternity ward for 1 day, then in the dispensary up to 21 days, then in the standard conditions for growth up to 180 days, then in the premises for rearing up to 360 days, and then in the premises for fattening up to 540 days of age.

To realize the bioresource potential of the meat qualities of the bull-calves in the technology of their growth and rearing, complex biopreparations were used developed from natural raw materials by the scientists of Chuvash State Agricultural Academy: Prevention-N-A (V.G. Semenov, F.P. Petryankin, V.A. Vasiliev and others) and Prevention-N-E (V.G. Semenov, D.A. Nikitin, V.A. Vasiliev and others). The animals of Test Group 1 were injected intramuscularly with Prevention-N-A at a dose of 3 ml on 2-3 and 7-9 days of life; the animals of Test Group 2 were injected intramuscularly with Prevention-N-E at the indicated dose and at the same time. To the control group no biopreparations were administered.

Prevention-N-A is a complex preparation to activate non-specific resistance of the organism and to realize the production potential of young animals. It is an aqueous suspension containing a polysaccharide complex of yeast cells of *Saccharomyces cerevisiae* immobilized in agar gel with the addition of a benzimidazole derivative and bactericidal preparation of aminoglycoside group.

Prevention-N-E is a complex preparation to stimulate non-specific resistance of the organism and to prevent diseases of farm animals. It is an aqueous suspension containing a polysaccharide complex of yeast cells *Saccharomyces cerevisiae* immobilized in agar gel with the addition of a benzimidazole derivative and an antibiotic group of macrolides.

Results. It is established that the microclimate in the premises for growth, rearing and fattening of bull-calves corresponded to zoo-hygienic norms.

The average daily rations for bull-calves in periods of growth up to 90 and 180 days, rearing up to 360 days and fattening up to 540 days provided the body's needs for energy and nutrients, mineral elements and vitamins in accordance with detailed feeding standards.

The use of Prevention-N-A and Prevention-N-E in the technology of bull-calves breeding stimulates their growth and development. Thus, by the end of the fattening period, the bull-calves of Test Groups 1 and 2 exceeded the control peers:

- by 20.8 and 16.8 kg in live weight,
- by 5.2 and 3.8 cm in the withers height,
- by 3.3 and 2.0 cm in the chest width behind the blades,
- by 2.3 and 1.9 cm in the depth of the chest;
- by 4.8 and 4.2 cm in the girth of the chest behind the blades,
- by 6.8 and 4.6 cm in the oblique length of the trunk;
- by 2.2 and 1.8 cm in the width of the hips;
- by 0.8 and 0.7 cm in the metacarpus, respectively (P <0.05-0.01).

The average daily growth rate of the animals of the test groups also turned out to be higher than in the control during all the periods of postnatal ontogenesis.

The index of leg length in the test groups decreased as they grew; the indices of being well-set and chest and pelvic indices, on the contrary, increased, while the bone index practically did not change.

The live weight in Test Group 1 and 2 (466.4 ± 3.03 kg and 462.4 ± 3.53 kg, respectively) was higher after fattening compared to the control group (445.6 ± 2.79 kg): by 20.8 kg (4.7%, P <0.001) and 16.8 kg (3.8%, P <0.01). The bull-calves of Test Group 1 (454.0 ± 3.51 kg) and 2 (449.6 ± 3.39 kg) had bigger pre-slaughter weight than the control group (430.7 ± 2.71 kg) by 23.3 kg (5.4%, P <0.001) and 18.9 kg (4.4%, P <0.01), respectively. The carcass weight of the bull-calves receiving intramuscular injections of Prevention-N-A was 16.5 kg (7.2%, P <0.001) bigger than in the control group, and with the use of Prevention-N-E it was 12.9 kg (5.6%, P <0.01) bigger. The slaughter weight of Test Group 1 was 18.0 kg (7.4%, P <0.001) bigger, of Test Group 2 it was 13.9 kg (5.7%, P <0.01) than in the control group. According to the slaughter outcome, Test Groups 1 and 2 also had an advantage in comparison with the control group by 1.1 and 0.8%, respectively.

Thus, against the background of immunoprophylaxis, biopreparations have been found to improve the slaughter qualities of bull-calves.

The data presented in Table. 1 shows that the bulls of Test Groups 1 and 2 exceeded their control peers by:

- weight of the chilled carcass - 16.1 and 11.9 kg (P <0.01);
- total meat outcome - 13.5 and 9.7 kg (P <0.05-0.01);
- total fat outcome - 1.5 and 1.0 kg (P <0.05-0.01);
- cartilage and tendons - 0.5 and 0.3 kg (P > 0.05);
- bones - 2.1 and 1.9 kg (P > 0.05), respectively.

The relative outcome of the tendons and bones in the experimental groups was, on the contrary, lower by 0.04 and 0.06%, respectively, and by 0.35 and 0.11% (P > 0.05) than in the control group.

Table 1 – Morphological composition of bull-calves carcasses

Parameter	Group		
	Control	Test 1	Test 2
Chilled carcass weight, kg	220.6±2.37	236.7±2.47**	232.5±2.55**
Meat weight, kg	172.1±2.22	185.6±2.31**	181.8±2.36*
Meat outcome, %	78.01	78.41	78.19
Fat weight, kg	12.0±0.32	13.5±0.22**	13.0±0.16*
Fat outcome, %	2.8	3.0	2.9
Cartilage and tendons weight, kg	8.3±0.12	8.8±0.25	8.6±0.19
Cartilage and tendons outcome, %	3.76	3.72	3.70
Bone weight, kg	40.2±0.75	42.3±0.66	42.1±0.71
Bone outcome, %	18.22	17.87	18.11
Meat outcome per 100 kg of pre-slaughter live weight	39.96±0.17	40.89±0.25*	40.45±0.23
Meat index	4.29±0.12	4.39±0.06	4.32±0.09

Legend: *P ≤ 0.05, **P ≤ 0.01.

The meat outcome per 100 kg of the pre-slaughter weight in Group 1 was 40.89 ± 0.25 kg (an increase by 0.93 kg or 2.3% (P <0.05)), while in Group 2 it was 40.45 ± 0.23 kg (an increase by 0.49 kg or 1.2% (P > 0.01)) than in the control group – 39.96 ± 0.17 kg. According to the meat index characterizing

the ratio of meat and bones, the bull-calves of Test Group 1 differed favorably – the aforementioned index was 4,39 – more than in the control group and Test Group 2 (0,10 and 0,07, accordingly).

Assessing the meat productivity, it is important to take into account not only the ratio of carcass tissues but also of the anatomical parts from which different types of meat are obtained. The analysis of the data received (table 2) indicates that the large carcass mass of test groups also determined high outcome of the most valuable cuts: brisket and sirloin – 6.1 and 4.0 kg ($P < 0.01-0.001$), rump – 2.6 and 1,7 kg ($P < 0,05-0,01$) and thick flank – 8,6 and 7,1 kg ($P < 0,001$), compared with the control group. In this case, the outcome of these cuts relative to the carcass mass of the bull-calves from Test Group 1 and 2 was higher by 0.7 and 0.3%, by 0.4 and 0.2%, by 1.4 and 1, 4% respectively, compared with the control group.

Table 2 – Weight and outcome of cuts from bull-calves carcasses

Parameter	Group		
	Control	Test 1	Test 2
Carcass weight, kg	220.6±2.37	236.7±2.47**	232.5±3.55**
cuts:			
neck, kg	23.8±0.12	23.4±0.24	23.5±0.22
%	10.8	9.9	10.1
chuck and blade, kg	41.0±0.22	40.2±0.20	40.4±0.19
%	18.6	17.0	17.4
brisket and sirloin, kg	61.8±0.66	67.9±0.51***	65.8±0.45**
%	28.0	28.7	28.3
rump, kg	23.2±0.40	25.8±0.48**	24.9±0.45*
%	10.5	10.9	10.7
thick flank, kg	70.8±0.85	79.4±0.80***	77.9±0.82***
%	32.1	33.5	33.5

Legend: * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$.

The carcasses of Test Groups 1 (27.8 kg) and 2 (26.7 kg), respectively, were characterized by the largest outcome of highest quality meat (table 3) - by 3.5 and 2.4 kg compared with the control group (24.3 kg, $P < 0.05-0.001$). At the same time, the relative outcome of highest quality beef in relation to the total meat weight was higher in the animals of the test groups – by 0.9 and 0.6% compared with the control group.

Table 3 – Meat quality composition of bull-calves carcasses

Parameter	Group		
	Control	Test 1	Test 2
Meat weight, kg	172.1±2.22	185.6±2.31**	181.8±2.36*
Highest quality meat weight, kg	24.3±0.73	27.8±0.72**	26.7±0.58*
Highest quality meat outcome, %	14.1	15.0	14.7
First-grade meat weight, kg	99.1±1.23	108.6±1.35***	105.6±1.29**
First-grade meat outcome, %	57.6	58.5	58.1
Second-grade meat weight, kg	48.7±0.62	49.2±0.60	49.5±0.59
Second-grade meat outcome, %	28.3	26.5	27.2

Legend: * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$.

From the culinary point of view, the meat quality composition of the individual anatomical parts of carcasses is of some interest (table 4).

The analysis of the data obtained indicates that the neck cut up to the seventh vertebra mainly consists of first- and second-grade meat. At the same time, the bull-calves of the test groups were inferior to the control group by the weight of the highest quality meat (0.1 kg) and first-grade meat (0.8 kg), yet the difference was unreliable. In case of chuck and blade cuts, the difference between all the three groups was insignificant ($P > 0.05$). In brisket and sirloin cuts, the largest highest quality meat outcome was in the test groups, the total outcome being 0.9 and 0.7 kg more, and the relative being 0.4 and 0.2% more.

Table 4 – Meat quality of cuts from bull-calves carcasses

Parameter	Group		
	Control	Test 1	Test 2
<i>Neck</i>			
Meat weight, kg	20,3±0,37	18,7±0,44	18,9±0,48
Highest quality meat weight, kg	1,9±0,13	1,8±0,14	1,8±0,17
Highest quality meat outcome, %	9,3	9,5	9,5
First-grade meat weight, kg	12,2±0,37	11,4±0,29	11,4±0,51
First-grade meat outcome, %	60,2	61,2	60,6
Second-grade meat weight, kg	6,2±0,25	5,5±0,32	5,7±0,25
Second-grade meat outcome, %	30,5	29,3	29,9
<i>Chuck and blade</i>			
Meat weight, kg	30,6±0,29	28,8±0,34	30,0±0,35
Highest quality meat weight, kg	4,1±0,19	4,0±0,16	4,1±0,10
Highest quality meat outcome, %	13,4	13,9	13,6
First-grade meat weight, kg	19,0±0,35	18,0±0,22	18,7±0,30
First-grade meat outcome, %	62,0	62,6	62,4
Second-grade meat weight, kg	7,5±0,22	6,8±0,20	7,2±0,25
Second-grade meat outcome, %	24,6	23,5	24,0
<i>Brisket and sirloin</i>			
Meat weight, kg	45,7±0,89	52,1±1,05**	51,0±0,84**
Highest quality meat weight, kg	4,6±0,19	5,5±0,16**	5,3±0,12*
Highest quality meat outcome, %	10,1	10,5	10,3
First-grade meat weight, kg	21,4±0,37	24,6±0,51***	24,0±0,42**
First-grade meat outcome, %	46,9	47,3	47,1
Second-grade meat weight, kg	19,7±0,34	22,0±0,47**	21,7±0,44**
Second-grade meat outcome, %	43,0	42,2	42,6
<i>Rump</i>			
Meat weight, kg	19,4±0,31	21,7±0,37**	20,5±0,32*
Highest quality meat weight, kg	3,0±0,11	3,5±0,17*	3,3±0,15
Highest quality meat outcome, %	15,7	16,1	15,9
First-grade meat weight, kg	11,8±0,24	13,2±0,25**	12,5±0,14*
First-grade meat outcome, %	60,8	61,0	60,9
Second-grade meat weight, kg	4,6±0,21	5,0±0,27	4,7±0,22
Second-grade meat outcome, %	23,5	22,9	23,2
<i>Thick flank</i>			
Meat weight, kg	56,1±0,97	64,3±0,94***	61,4±0,81**
Highest quality meat weight, kg	10,7±0,18	13,0±0,22***	12,2±0,25**
Highest quality meat outcome, %	19,1	20,2	19,9
First-grade meat weight, kg	34,7±0,68	41,4±0,75***	39,0±0,71**
First-grade meat outcome, %	61,8	64,4	63,5
Second-grade meat weight, kg	10,7±0,12	9,9±0,17	10,2±0,19
Second-grade meat outcome, %	19,1	15,4	16,6

The amount of highest grade meat in the rump cuts in Test Group 1 was 0.5 and 0.2 kg more than in the control and Test Group 2, respectively; the relative outcome was also 0.4 and 0.2% more, respectively. The largest and most valuable cut of the carcass is the thick flank, as it gives the highest outcome of highest quality meat; in Test Groups 1 and 2 this amount was 2.3 and 1.5 kg more ($P < 0.01-0.001$) than in the control group. At the same time, the relative outcome of the highest quality meat was 19.1% in the control group, 20.2% in Test Group 1 and 19.9% in Test Group 2.

Thus, brisket and sirloin, rump and thick flank cuts in the test groups were characterized by the highest outcome of highest quality meat in comparison with the control group.

The results of beef quality assessment by organoleptic, biochemical and spectrometric indicators are presented in table 5.

Table 5 – Assessment of beef quality

Parameter	Group		
	Control	Test 1	Test 2
Organoleptic:			
surface appearance and color	the meat samples are covered with a dried-up crust of pale pink color		
muscles on the cut	slightly moist, do not leave a wet spot on the filter paper; light red in color		
consistency	meat is dense, elastic on the cut; the dimple formed with a finger is quickly aligned		
smell	specific, typical of fresh beef		
surface fat	has a yellowish color; the consistency is hard, crumbles when pressed		
tendons condition	elastic, dense; joints surface is smooth and shiny		
broth transparency and smell	transparent, fragrant, large drops of fat on the surface		
Biochemical:			
pH (5,6 – 6,2)	5,96±0,03	5,98±0,01	5,94±0,02
amino-ammonium nitrogen, mg (not more than 1,26in 10 ml extracts from fresh meat)	1,20±0,01	1,17±0,01	1,18±0,01
formalin test	negative		
peroxidase test	positive		
copper sulphate test	negative		
Spectrometric, mg/kg			
lead (not more than 0,5)	0,08±0,01	0,08±0,01	0,07±0,01
cadmium (not more than 0,05)	not found	same as control	same as control
arsenic (not more than 0,1)	not found	same as control	same as control
copper (not more than 5,0)	0,81±0,04	0,84±0,02	0,83±0,03
zinc (not more than 70)	27,5±0,24	27,8±0,16	28,2±0,21
mercury (not more than 0,03)	not found	same as control	same as control

The beef had a dry crust and a pale pink color. The cut was uneven, soaked with blood more intensively than in other parts of the carcass; there was no blood in the muscles and blood vessels. Small vessels under the pleura and peritoneum were not visible. The lymph node cuts were light gray in color. Meat was dense, elastic; when pressing a finger on its surface a dimple was formed which quickly disappeared. The muscles on the incision were slightly moistened and did not leave a moist spot on the filter paper, had a light red color. The broth prepared from the meat was transparent, fragrant, with a small accumulation of large drops of fat on its surface.

It was found that the beef pH in the test experimental groups varied within a narrow range and amounted to 5.96 ± 0.03 , 5.98 ± 0.01 and 5.94 ± 0.02 in the control, Test Group 1 and 2, respectively. The content of amino-ammonium nitrogen in the beef samples was 1.20 ± 0.01 mg (control group), 1.17 ± 0.01 (Test Group 1) and 1.18 ± 0.01 mg (Test Group 2) – in other words, within the norm. In the studied samples the reaction with formalin turned out to be negative, with peroxidase - positive, and with copper sulphate - negative.

In lead concentration, the samples were practically indistinguishable – it averaged 0.08 ± 0.01 mg/kg in the control group, 0.08 ± 0.01 in Test Group 1 and 0.07 ± 0.01 mg/kg in Test Group 2. Such toxic elements as cadmium, arsenic and mercury in beef samples were not detected. The level of copper and zinc in the meat samples was within the permissible limits: 0.81 ± 0.04 and 27.5 ± 0.24 mg/kg in the control group, 0.84 ± 0.02 and 27.8 ± 0.16 mg/kg in Test Group 1 and 0.83 ± 0.03 and 28.2 ± 0.21 mg/kg in Test Group 2. Consequently, by spectrometric parameters the meat of the bull-calves of the experimental groups practically did not differ from the control data.

Thus, the veterinary and sanitary examination found that organoleptic, biochemical and spectrometric indicators of meat of bull-calves grown on the background of intramuscular injection of Prevention-N-A and Prevention-N-E did not differ from those in the control group and met the requirements of the

Technical Regulations of the Customs Union 'On Food Safety' TR CU 021/2011 and the Technical Regulations of the Customs Union 'On the Safety of Meat and Meat Products' TR CU 034/2013, which indicates the safety of the test preparations and good quality of meat carcasses.

Based on the analysis of the age-related dynamics of the clinical and physiological state, it was established that body temperature, pulse rate and respiratory movements in the calves of the compared groups during the periods of growth, rearing and fattening were within the physiological norms, and the difference in them was insignificant ($P > 0, 05$).

Prevention-N-A and Prevention-N-E tested in experiments on Black Motley breed activate the production of red blood cells and increase the concentration of hemoglobin in the blood, that is, improve haemopoiesis, but do not affect leucopoiesis.

Against the background of immunoprophylaxis of experimental bull-calves, cellular and humoral factors of non-specific protection are activated, which is especially important in the early period of postnatal ontogeny. On the 30th day of the growth period, the bulls of Test Groups 1 and 2 exceeded the control peers by 4.8 and 4.2% in phagocytic activity of the leukocytes, in phagocytic index by 1.1 and 0.8, in lysozyme activity in plasma by 2.1 and 1.5%, in bactericidal activity in serum by 6.3 and 5.5%, and in the concentration of immunoglobulins in blood serum by 3.1 and 2.1 mg/ml.

The phagocytic activity of neutrophils segmented leukocytes towards *Staphylococcus aureus* in the bull-calves in the three groups gradually increased as they grew and developed - from 31.0 ± 1.26 to $64.6 \pm 1.44\%$, from 30.6 ± 1.21 to $71.6 \pm 1.47\%$ and from 30.8 ± 1.02 to $70.0 \pm 1.41\%$, respectively (control, Test Group 1, Test Group 2). The most pronounced activity of this cellular factor of non-specific resistance of the organism was observed in 30, 90, 180, 360 and 540-day-old animals of Test Group 1 (intramuscular injections of Prevention-N-A) in comparison with the control data for 4.8, 6.8, 7.0, 6.9, and 7.0% ($P < 0.05-0.01$). This activity of leukocytes in the animals of Test Group 2 (intramuscular injections of Prevention-N-E) was also significantly higher than in the control group, starting from their 30-day-old age and till the fattening period: in 30-day-old bull-calves by 4.2%, 90-day - 5.6%, 180-day - 5.6%, 360-day - 5.4% and 540-day-old - 5.4% ($P < 0,05$).

The phagocytic index in the animals of the three groups increased from the 1st to the 5th day of studies from 2.3 ± 0.12 to 9.2 ± 0.37 , from 2.1 ± 0.10 to 10.2 ± 0.41 and from 2.5 ± 0.22 to 9.4 ± 0.40 , respectively. It should be noted that the average number of bacteria in one phagocyte in the animals of Test Group 1 was significantly higher than in the control group on the 30th, 90th and 180th days of the growth period (by 16.6%, 16.7 and 13.7%, respectively) and on the 360th day of the rearing period by 17.5% ($P < 0.05$). The phagocytic index in the bulls of Test Group 2 was also higher in comparison with the control group, yet the difference proved to be reliable only on the 30th and 90th day - by 12.2 and 12.5% ($P < 0.05$).

The lysozyme activity in the blood plasma in the control group and in Test Group 2 increased consecutively during the period of growth - from the 1st to the 90th day, respectively by 7.1 ± 0.36 to $20.3 \pm 0.55\%$, and from 7.3 ± 0.35 to $22.6 \pm 0.42\%$. However, on the 180th day of the indicated period it decreased to 20.0 ± 0.41 and $22.2 \pm 0.58\%$, and subsequently during the periods of rearing and fattening it steadily increased, reaching the peak at the end of the fattening period - 24.4 ± 0.51 and $26.4 \pm 0.40\%$. The muramidase activity of this enzyme in Test Group 1 was continuously increased during the experiment from 7.1 ± 0.33 to $26.5 \pm 0.39\%$. This indicator of the humoral link of the non-specific resistance of the organism in the animals of Test Groups 1 and 2 was significantly higher than in the control group, starting from the 30-day of age till slaughter: in 30-day-old animals by 2.1 and 1.5%, in 90-day-old by 3.4 and 2.3%, in 180-day-old animals by 3.9 and 2.2%, in 360-day-old by 2.3 and 1.8%, and in 540-day-old by 2.1 and 2.0% ($P < 0.05-0.001$), respectively.

It was found that bactericidal activity of blood serum of tested animals tended to increase as they grew and developed from the 1st to the 540th day: in the control group the increase was from 33.6 ± 1.07 to $60.0 \pm 0.85\%$, in Test Group 1 from 33.3 ± 1.19 to $63.3 \pm 0.96\%$, and in Test Group 2 from 33.4 ± 1.14 to $61.9 \pm 1.16\%$. Here, the bactericidal activity of blood serum in Test Group 1 was higher than the control data for all the periods of research: on the 30th day by 6.3%, on the 90th day - 6.1%, on the 180th day - 4.6%, on the 360th day - 3.8% and on the 540th day by 3.3% ($P < 0.05-0.01$). It should be noted that this activity in Test Group 2 also had a higher rate than that in the control group, especially during the growth period. Thus, the 30-day-old bull-calves of this group exceeded their control peers by this factor of the

humoral link of the non-specific resistance by 5.5%, 90-day-old bull-calves by 5.5% and 180-day-olds by 5.6% ($P < 0, 05-0.01$).

It was established that the concentration of immunoglobulins in the blood serum of the bull-calves of the test groups increased as they grew and developed: in the control group, from 11.6 ± 0.68 to 28.1 ± 0.93 mg/ml, in Test Group 1 from 11.5 ± 0.64 to 32.1 ± 0.76 mg/ml and in Test Group 2 from 11.4 ± 0.70 to 31.8 ± 0.72 mg/ml. The level of indicated immunocompetent factor of blood serum in the test groups was significantly higher – by 3.1 and 2.1 mg/ml, 3.6 and 2.0 mg/ml, 4.1 and 2.9 mg/ml, 3.9 and 3.1 mg/ml and 4.0 and 3.7 mg/ml (i.e., 19.2 and 13.0%, 16.3 and 9.0%, 15.6 and 11.1%, 13.9 and 11.1% and 14.2 and 13.2%) after 30, 90, 180, 360 and 540 days after the experiments, compared with the control group ($P < 0.050.01$).

Based on the analysis of immunological studies, it was established that bull-calves receiving biopreparations and reared and fattened in standard conditions had higher rates of cellular and humoral non-specific defense of the organism. Moreover, the immunostimulating effect of Prevention-N-A was more pronounced than Prevention-N-E.

Discussion. At the present stage of cattle breeding development to ensure reliable health protection and realization of the bioresource potential and the meat qualities of bull-calves, it becomes necessary to activate the non-specific protective factors of the organism against the technological and environmental factors of the habitat during periods of growth, rearing and fattening by biological preparations which are characterized by high bioavailability and harmlessness to the organism.

The pharmaceutical market offers a wide range of diverse products, many of chemical origin and thus of low bioavailability. In addition, the previously proposed drugs only work on individual factors of non-specific resistance, which does not fully ensure the activation of the immune system. In secondary immune deficiencies, pathogenic microorganisms play an important role in the development of the disease, therefore, in the treatment of animals antibiotics that can have an immunosuppressive effect are used [8, 10, 14, 16, 23, 25, 27, 28, 31, 33].

The authors of the present study believe it is most expedient to use immunostimulants made of natural raw materials with antibacterial drugs. When combined, a double strike is applied to the pathogen: the antibacterial drug suppresses the functional activity of the pathogen, increasing its sensitivity to phagocytosis, and the immunostimulant activates the phagocyte, increasing its ability to neutralize the pathogen.

In view of the foregoing, a scientific study was devoted to the realization of the bioresource potential of meat qualities of Black Motley bull-calves by directed correction of postnatal formation and development of non-specific resistance of the organism with the help of biopreparations Prevention-N-A and Prevention-N-E in connection with hygienic conditions of maintenance and feeding.

The dynamics of the live mass, both absolute and relative, gives an accurate prediction about the development of meat productivity of an animal, both in its lifetime and after slaughter [32]. In conditions of two intramuscular injections of Prevention-N-A and Prevention-N-E on the 2nd-3rd and 7th-9th day in a dose of 3 ml growth and development boost was established. By the end of the growth period, 180-day-old bull-calves of the test groups outperformed the control peers by 7.2 and 4.8 kg of live weight, by 14.6 and 12.0 kg during rearing and by 20.8 and 16.8 kg after fattening, respectively ($P < 0.05-0.001$).

The classics specialist of Russian zootechnics Bogdanov E.A., Kuleshov N.N., Ivanov M.F., Gol-dobin M.I. and others pointed out that only constitutionally strong animals meet economic and biological requirements. The data obtained and the analysis of the external-constitutional features of the bull-calves in the dynamics show that the animals of the test groups exceeded the control peers both in height and girth. For example, at the end of the fattening period the parameters were as follows: withers height by 5.2 and 3.8 cm, chest width behind the blades by 3.3 and 2.0 cm, chest depth by 2.3 and 1.9 cm, chest girth behind the blades by 4.8 and 4.2 cm, oblique length of the trunk by 6.8 and 4.6 cm, hips width by 2.2 and 1.8 cm, and in metacarpus by 0.8 and 0.7 cm, respectively.

In order to assess the meat productivity and quality, control bull-calves were slaughtered at the age of 18 months (5 animals from each group). It was found that the bull-calves of Test Groups 1 and 2 exceeded the control peers by the weight of the chilled carcass by 16.1 and 11.9 kg ($P < 0.01$), the total meat outcome by 13.5 and 9.7 kg ($P < 0,05-0,01$), fat by 1.5 and 1.0 kg ($P < 0,05-0,01$), cartilage and tendons by 0.5 and 0.3 kg ($P > 0.05$), bones by 2.1 and 1.9 kg ($P > 0.05$), respectively. The relative outcome of tendons and

bones in test groups was, on the contrary, lower by 0.04 and 0.06%, respectively, and by 0.35 and 0.11% ($P > 0.05$) than in the control group. Thus, full-bodied carcasses were obtained from test groups; nevertheless, the bull-calves of Test Group 1 were leading along all the indicators.

Nutritional value, taste and culinary benefits of various natural anatomical parts of the carcass are not the same. In the present experiment, the larger weight of carcasses of the bull-calves of the test groups determined the high outcome of the most valuable cuts: brisket and sirloin by 6.1 and 4.0 kg ($P < 0.01-0.001$), rump by 2.6 and 1.7 kg ($P < 0.05-0.01$) and thick flank by 8.6 and 7.1 kg ($P < 0.001$) compared with the control group.

The composition of meat largely determines its further use by meat processing plants, and the range of meat products. The largest outcome of highest quality meat was from the carcasses of the test groups (3.5 and 2.4 kg, respectively) in comparison with the control group ($P < 0.05-0.001$). In the carcass, the largest and most valuable cut is the thick flank, as it gives the highest outcome of premium quality meat. The amount of such meat in the test groups was 2.3 and 1.5 kg larger ($P < 0.01-0.001$) than in the control group.

The quality of meat products is determined by its chemical composition and biological wholesomeness, which in turn is determined by the conformity of the product to the needs of the human body and the guaranteed harmlessness of its use in accordance with physiological standards [12]. As a result of the veterinary and sanitary examination, the compliance of beef with the requirements of the Technical Regulations of the Customs Union 'On Food Safety' TR CU 021/2011 and the Technical Regulations of the Customs Union 'On the Safety of Meat and Meat Products' TR CU 034/2013 was established, indicating the good quality of meat carcasses.

Prevention-N-A and Prevention-N-E tested on Black Motley bull-calves increased the number of erythrocytes and the hemoglobin concentration in blood, that is, improved haemopoiesis, and also activated cellular and humoral factors of non-specific resistance.

Conclusions. Under the influence of Prevention-N-A and Prevention-N-E the growth and development of Black Motley bull-calves is accelerated during the periods of growth, rearing and fattening, which causes their higher slaughter and meat qualities and, as a consequence, the outcome of valuable cuts – brisket and sirloin, rump and thick flank, and also of highest quality and first-grade beef. It was experimentally proved that the realization of the bioresource potential is caused by the activation of haemopoiesis, cellular and humoral factors of the non-specific resistance with the help of biological preparations, with a more pronounced corresponding effect of Prevention-N-A.

1. Using Prevention-N-A and Prevention-N-E in the technology of growing bull-calves twice on the 2nd-3rd and 7th-9th day in the dose of 3 ml stimulates their growth and development.

By the end of the fattening period the bull-calves of the test groups exceeded the control peers by 20.8 and 16.8 kg in live weight, by 5.2 and 3.8 cm at the withers, by 3.3 and 2.0 cm in chest width behind the blades, by 2.3 and 1.9 cm in chest depth, by 4.8 and 4.2 cm in chest girth behind the blades, by 6.8 and 4.6 cm in the oblique length of the trunk, by 2.2 and 1.8 cm in hips width and by 0.8 and 0.7 cm in metacarpus, respectively ($P < 0.05-0.01$). The average daily growth of the animals of the test groups also turned out to be higher than in the control during all the periods of postnatal ontogeny.

Leg length index of the animals of the test groups decreased as they grew, the indices of stretch, breach, chest and pelvic, on the contrary, increased, and the bone index practically did not change.

2. Against the background of the application of biological preparations, the fattening and slaughtering qualities of the bull-calves improve.

An increase in the pre-slaughter live weight of animals of the test groups by 23.3 and 18.9 kg was established, the weight of the carcass was increased by 16.5 and 12.9 kg, the slaughter outcome by 1.1 and 0.8%, the total meat outcome by 13.5 and 9.7 kg, internal fat by 1.5 and 1.0 kg and meat index by 0.10 and 0.07, respectively ($P < 0.05-0.01$). The outcome of the most valuable cuts also increased: brisket and sirloin by 6.1 and 4.0 kg ($P < 0.01-0.001$), rump by 2.6 and 1.7 kg ($P < 0.05-0.01$) and thick flank by 8.6 and 7.1 kg ($P < 0.001$) than in the control group.

3. Including Prevention-N-A and Prevention-N-E in the technology of growing bull-calves promotes the improvement of meat qualities.

The largest outcome of highest quality meat was from the carcasses of the bull-calves of Test Group 1 (27.8 ± 0.72 kg) and 2 (26.7 ± 0.58 kg), respectively by 3.5 and 2.4 kg, as compared with the control

group (24.3 ± 0.3 kg), and also of their cuts: brisket and sirloin by 0.9 and 0.7 kg, rump by 0.5 and 0.3 kg, thick flank by 2.3 and 1.5 kg ($P < 0.05-0.001$).

The beef conformed to the requirements of the Technical Regulations of the Customs Union 'On Food Safety' TR CU 021/2011 and the Technical Regulations of the Customs Union 'On the Safety of Meat and Meat Products' TR CU 034/2013.

4. The use of Prevention-N-A and Prevention-N-E in the technology of growing, rearing and fattening bull-calves does not affect the clinical and physiological state of the organism.

5. Prevention-N-A and Prevention-N-E tested on Black Motley bull-calves activate the production of erythrocytes and increase the concentration of hemoglobin in the blood, that is, improve haemopoiesis, but do not affect leucopoiesis.

6. Against the background of immunoprophylaxis of experimental bull-calves, cellular and humoral factors of non-specific resistance are activated, which is especially important in the early period of postnatal ontogeny. On the 30th day of the growth period the animals of the test groups exceeded the control peers by 4.8 and 4.2% in phagocytic activity of the leukocytes, in phagocytic index by 1.1 and 0.8, in lysozyme activity of the plasma by 2.1 and 1.5%, in bactericidal activity of serum by 6.3 and 5.5%, and in the concentration of immunoglobulins in blood serum by 3.1 and 2.1 mg/ml.

Recommendations. To implement the bioresource potential of meat qualities of Black Motley bull-calves the authors recommend the use of complex biopreparations Prevention-N-A and Prevention-N-E, which are immunostimulants based on the polysaccharide complex of yeast cells *Saccharomyces cerevisiae* in combination with bactericidal preparations of aminoglycoside and natural macrolides:

1) intramuscularly injecting Prevention-N-A twice daily for newborn calves for on the 2nd-3rd and 7th-9th day in the dose of 3 ml;

2) intramuscularly injecting Prevention-N-E twice on the 2nd-3rd and 7th-9th day in the dose of 3 ml.

The proposed biopreparations contribute to the realization of the bioresource potential of the meat qualities of bull-calves due to the activation of the protective-adaptive functions of the organism against environmental and technological factors and the selective mobilization of the hematologic profile and cellular and humoral factors of non-specific resistance, with a more pronounced corresponding effect of Prevention-N-A.

REFERENCES

[1] Amerkhanov Kh.A. Scientific ensurance of competitiveness of dairy cattle breeding [Nauchnoye obespecheniye konkurentosposobnosti molochnogo skotovodstva] / Kh.A. Amerkhanov, N.I. Strekozov // Dairy and meat cattle breeding. M., 2012. N 1. P. 2-5.

[2] Basonov O.A. Black-and-white cattle of Nizhny Novgorod selection: Monograph. [Cherno-pestryy skot nizhegorodskoy selektsii] / O.A. Basonov, M.E. Taigunov, A.V. Katkov, A.V. Shishkin // Nizhny Novgorod: Quartz, 2016. 260 p.

[3] Vasiliev V.A. The use of biological products in the technology of growing, rearing and fattening of bull-calves [Ispol'zovaniye biopreparatov v tekhnologii vyrashchivaniya, dorashchivaniya i otkorma bychkov] / V.A. Vasiliev, V.G. Semenov // Youth and Innovation: materials of Russian scientific-practical. conf. of young scientists, graduates and students. Cheboksary, 2017. P. 68-70.

[4] Gizatullin, R.S. adaptive resource-saving technology of beef production in meat cattle-breeding: Monograph. / R.S. Gizatullin, T.A. Sedykh [Adaptivnaya resursoberegayushchaya tekhnologiya proizvodstva govyadiny v myasnom skotovodstve] // Saarbrücken, 2016. 119 p.

[5] Gilmiyarov A.L. Slaughter qualities of young Black Motley breed and its half-breed hybrids with Aubrac [Uboynnye kachestva molodnyaka cherno-pestroy porody i yego polukrovnykh pomesey s porodoy obrak] / A.L. Gilmiyarov, H.H. Tagirov, I.V. Mironova // Bulletin of Bashkir State Agrarian University. Ufa, 2010. N 3. P. 15-19.

[6] Goldobin M.I. Beef production reserves / Use of replacements for fattening [Rezervy proizvodstva govyadiny / Ispol'zovaniye sverkhremontnykh telok dlya otkorma] / M.I. Goldobin, A.G. Grigoriev, R.M. Aizatov // Zootechnics. M., 1994. N 11. P. 26-27.

[7] Gorlov I.F. Efficiency of beef production with the use of new anti-stress lactulose-containing preparations: Recommendations. [Effektivnost proizvodstva govyadiny pri ispol'zovanii novykh antistressovykh laktulozoderzhashchikh preparatov] / I.F. Gorlov, M.I. Slozhenkina, A.A. Zakurdaeva, A.V. Randelin, D.A. Mosolova, A.S. Miroshnik. Volgograd, 2017. 19 p.

[8] Dementiev E.P. Modern problems of zoohygiene and ways to solve them / E.P. Dementiev, V.G. Tyurin [Sovremennyye problemy zoogigiyeny i puti ikh resheniya] // 30 years of the department of zoohygiene, epizootology and the basis of veterinary medicine. Ufa, 2000. P. 24-28.

[9] Dzhaparidze T.G. Creating a branch of beef cattle breeding [Sozdat otrasl myasnogo skotovodstva] / T.G. Dzhaparidze // Leading zootechnician. M., 2008. N 8. P. 39-41.

[10] Donnik I.M. Acute respiratory diseases of cattle and prevention problems in modern industrial production conditions [Ostryye respiratornyye zabolvaniya krupnogo rogatogo skota i problemy profilaktiki v sovremennykh usloviyakh promysh-

lennogo proizvodstva] / Donnik, O.G. Petrova, S.A. Markovskaya // Agrarian Bulletin of the Urals. Ekaterinburg, 2013. N 10(116). P. 25-27.

[11] Dunin I.M. Selection and technological aspects of the development of dairy cattle breeding in Russia [Selektsionno-tekhnologicheskiye aspekty razvitiya molochnogo skotovodstva Rossii] / I.M. Dunin, H.A. Amerkhanov // Zootechnics. M., 2017. N 6. C. 2-8.

[12] Iskhakov R.G. Growth and meat productivity of purebred and crossed bull-calves of different genotypes [Rost i myasnaya produktivnost chistoporodnykh i pomesnykh bychkov razlichnykh genotipov] / R.G. Iskhakov, V.I. Levakhin, R.M. Galiev // Bulletin of beef cattle breeding. Orenburg, 2006. N 59 (1). P. 119-121.

[13] Labinov V.V. Modernization of Black Motley breed of cattle in Russia on the basis of using the Holstein gene pool [Modernizatsiya cherno-pestroy porody krupnogo rogatogo skota v Rossii na osnove ispolzovaniya genofonda golshinov] / V.V. Labinov, P.N. Prokhorenko // Dairy and beef cattle. M., 2015. N 1. P. 2-7.

[14] Mudarisov R.M. Exterior-constitutional and economic-biological features of Holstein cows [Eksteryerno-konstitutsionalnyye i khozyaystvenno-biologicheskiye osobennosti korov golshinskoy porody] / R.M. Mudarisov, G.R. Ahmetzyanov, V.G. Semenov // Food security and sustainable development of agroindustrial complex: mat. Intern. scientific-practical. conf. Cheboksary, 2015. P. 449-454.

[15] Musayev F.A. Innovative technologies in the production of beef: Monograph. [Innovatsionnyye tekhnologii v proizvodstve govyadiny]. Musaev, N.I. Morozova. Ryazan, 2014. 160 p.

[16] Musayev F.A. Biologically active additives: application, safety, quality assessment: Monograph. [Biologicheski aktivnyye dobavki: primeneniye, bezopasnost', otsenka kachestva] / F.A. Musaev, O.A. Zakharova. Ryazan, 2016. 201 p.

[17] Petryankin F.P. Polysaccharides as immunity stimulants [Polisakharidy kak stimulyatory immuniteta] / F.P. Petryankin // The role of the higher school in the project 'Living thinking as strategy of Chuvashia': materials of Intern. scientific-practical. Conf. Cheboksary, 2010. P. 160-164.

[18] Petryankin F.P. Immunostimulants in the practice of veterinary medicine: Monograph. [Immunostimulyatory v praktike veterinarnoy meditsiny] / F.P. Petryankin, V.G. Semenov, N.G. Ivanov. Cheboksary: The New Time, 2015. 272 p.

[19] Semenov V.G. Cultivation of calves under different regimes of adaptive technology with application of domestic biopreparations [Vyrashchivaniye telyat pri raznykh rezhimakh adaptivnoy tekhnologii s primeneniyyem otechestvennykh biopreparatov] / V.G. Semenov, A.F. Kuznetsov, D.A. Nikitin, V.A. Vasiliev // Questions of legal regulation in veterinary medicine. SPb., 2016. N 4. P. 139-141.

[20] Semenov V.G. Realization of reproductive qualities of cows and productive potential of calves by biological preparations [Realizatsiya vosproizvoditel'nykh kachestv korov i produktivnogo potentsiala telyat biopreparatami] / V.G. Semenov, D.A. Nikitin, N.I. Gerasimova // Proceedings of the International Academy of Agrarian Education. SPb., 2017. N 33. P. 172-175.

[21] Semenov V.G. Nonspecific resistance of the animal organism to different stress factors [Nespetsificheskaya ustoychivost organizma zhivotnykh k stress-faktoram raznykh sil]. V.G. Semenov, D.A. Nikitin, A.V. Volkov, K.V. Zakharova // Ecology of the native land: problems and solutions: mat. XII Russian scientific-practical. conf. with intern. participation in the Forum 'EcoKirov-2017'. Kirov, 2017. P. 233-237.

[22] Semenov V.G. Realization of bioresource potential of black-and-white cattle by biological preparations [Realizatsiya bioresursnogo potentsiala cherno-pestrogo skota biopreparatami] / V.G. Semenov, D.A. Nikitin, N.I. Gerasimova, V.A. Vasiliev // Bulletin of Chuvash State Agricultural Academy. Cheboksary, 2017. N 1(1). P. 37-43.

[23] Smirnov A.M. Protection of agricultural animals against diseases is an important factor in improving livestock efficiency [Zashchita selskokhozyaystvennykh zhivotnykh ot bolezney – vazhnyy faktor povysheniya effektivnosti zhivotnovodstva] / A.M. Smirnov // Veterinary Medicine and Feeding. M., 2012. N 3. P. 4-12.

[24] Smirnova M.F. State and ways of increasing the production of beef in dairy cattle breeding in Leningrad Region [Sostoyaniye i puti uvelicheniya proizvodstva govyadiny v molochnom skotovodstve Leningradskoy oblasti] / M.F. Smirnova, V.V. Smirnova // Proceedings of St. Petersburg State Agrarian University. SPb., 2017. N 2(47). P. 231-235.

[25] Sofronov V.G. The use of the biological stimulator 'Univetselp' to correct metabolism in cows and prevent dyspepsia in calves [Primeneniye biologicheskogo stimulyatora 'Univetselp' dlya korrektsii obmena veshchestv u korov i profilaktiki dispetsii telyat] / V.G. Sofronov, U.Z. Ibragimov, R.Kh. Gadzaonov, F.N. Chekhodaridi // Sci. papers of Bauman Kazan State Academy of veterinary medicine. Kazan, 2011. Vol. 205. P. 200-206.

[26] Strekozov N.I. Evaluation of dairy breeds by reproductive and adaptive abilities [Otsenka molochnykh porod po vosproizvoditel'nykh i adaptatsionnykh sposobnostyam] / N.I. Strekozov, N.V. Sivkin, V.I. Chinarov, O.V. Bautina // Zootechnics. M., 2017. N 7. P. 2-6.

[27] Topuria L.Yu. Effect of drugs of natural origin on the reproductive ability and immune status of cows [Vliyaniye preparatov prirodnoy proiskhozhdeniya na vosproizvoditel'nyuyu sposobnost i immunnyy status korov]. L.Yu. Topuria, G.M. Topuria // Bulletin of Altai State Agricultural University. Barnaul, 2007. N 5. P. 52-55.

[28] Khakimov I.N. The use of hormonal drugs to intensify the reproduction of herds [Ispolzovaniye gormonalnykh preparatov dlya intensivatsii vosproizvodstva stad] / I.N. Khakimov, R.M. Mudarisov // Integration of science and practice as a mechanism of effective development of the agroindustrial complex: mat. Intern. scientific-practical. conf. in the framework of the XIII International exhibition 'AgroComplex-2013'. Ufa, 2013. P. 282-285.

[29] Khusainov I.I. The main directions of development of beef production technologies [Osnovnyye napravleniya razvitiya tekhnologiy proizvodstva govyadiny] / I.I. Khusainov, I.Yu. Morozov // Effective livestock breeding. Krasnodar, 2015. N 8(117). P. 36-40.

[30] Shevkhuzhev A.F. The quality of meat obtained with different technologies of growing bulls [Kachestvo myasa, poluchennogo pri raznykh tekhnologiyakh vyrashchivaniya bychkov] / A.F. Shevkhuzhev, RA Ulimbasheva // Bulletin of Altai State Agricultural University. Barnaul, 2015. N 3(125). P. 140-143.

[31] Shukanov A.A. Characteristics of changes in the clinical and physiological state, productivity and quality of meat in bull-calves when prescribing domestic preparations [Kharakteristika izmeneniy kliniko-fiziologicheskogo sostoyaniya, produktivnosti i kachestva myasa u bychkov pri naznachenii otechestvennykh biopreparatov] / G.A. Yakovlev, A.A. Shukanov, I.F. Kabirov // Veterinary Physician. Kazan, 2013. N 3. P. 54-56.

[32] Yuberova A.A. Peculiarities of the formation of meat qualities of Black Motley bull-calves in the Michurin agricultural production cooperative [Osobennosti formirovaniya myasnykh kachestv bychkov cherno-pestroy porody v SKHPK im. Michurina] / A.A. Yuberova, O.A. Krasnova, O.S. Starostina // Bulletin of Izhevsk State Agricultural Academy. Izhevsk, 2012. N 2(31). P. 41-43.

[33] Yakimov A.V. Prospects of using new feed additives in cattle rations [Perspektivy ispol'zovaniya novykh kormovykh dobavok v ratsionakh krupnogo rogatogo skota] / A.V. Yakimov, M.G. Ziatdinov, V.V. Gromakov // Bulletin of Ulyanovsk State Agricultural Academy. Ulyanovsk, 2014. N 3(274). P. 96-98.

[34] Yakimov A.V. Technologies for the livestock production using new feed additives [Tekhnologii proizvodstva zhivotnovodcheskoy produktsii s ispolzovaniyem novykh kormovykh dobavok] / A.V. Yakimov, F.Zh. Mudarisov, V.V. Salakhov // Bulletin of Ulyanovsk State Agricultural Academy. Ulyanovsk, 2016. N 3(35). P. 165-169.

[35] Semenov V.G. Health and productivity of bulls under different modes of cultivation, rearing and fattening / V.G. Semenov, R.M. Mudarisov, V.A. Vasilyev // 'European conference on innovations in technical and natural sciences'. Proceedings of the 1st International scientific conference. Vienna, Austria 2014. Section 9. Agricultural sciences. P. 176-181.

**В. Г. Семенов¹, Д. А. Баймұқанов², Н. И. Косяев¹, Р. М. Мударисов³,
Н. И. Морозова⁴, Ф. А. Мусаев⁴, Д. А. Никитин¹, М. Б. Қалмағамбетов²**

¹Чуваш мемлекеттік ауыл шаруашылық академиясы, Чебоксар, Шуас Республикасы, Ресей,

²Қазақ мал шаруашылығы және азық өндіру ғылыми-зерттеу институты, Алматы, Қазақстан,

³Башқұрт мемлекеттік аграрлық университеті, Уфа, Башқұрт Республикасы, Ресей,

⁴П. А. Костычев атындағы Рязань мемлекеттік агротехнологиялық университеті,
Рязань, Рязань облысы, Ресей

PREVENTION СЕРИЯЛЫ БИОПРЕПАРАТТАРДЫ ҚОЛДАНУ АЯСЫНДА БҰҚАЛАРДЫҢ ӨСУІ, ДАМУЫ ЖӘНЕ ЕТТІЛІК САПАЛАРЫ

Аннотация. Алғаш рет кешенді зерттеулер негізінде қара-ала тұқымды бұқалардың еттілік сапасының биоресурстық әлеуетін іске асыру үшін ойлап табылған Prevention-N-A және Prevention-N-E биопрепараттарын қолданудың зоотехникалық мақсатқа сәйкестігі ғылыми негізделгендігі және тәжірибие жүзінде дәлелденді. Биопрепараттарды қолдану аясында бұқаларды өсіру, өсіріп жеткізу және бордақылау кезінде өсу белсенділігі мен дамуы орнатылды, бұл өз кезегінде ұшаның сойыстық және еттілік сапаларын ас жоғарылауына алып келді, соның нәтижесінде, құнды кесектер шықты: тексерудегіге қарағанда арқатөстік – 6,1 және 4,0 кг-ға ($P<0,01-0,001$), белдік – 2,6 және 1,7 кг-ға ($P<0,05-0,01$) және жамбастық – 8,6 және 7,1 кг-ға ($P<0,001$) ұлғайды. Жоғары сортты жұмсақ еттер 1-ші ($27,8\pm 0,72$ кг) және 2-ші ($26,7\pm 0,58$ кг) тәжірибелік топтардағы бұқа ұшаларында байқалды, 3,5 және 2,4 кг-ға сәйкес тексерудегімен салыстырғанда ($24,3\pm 0,73$ кг), және де олардың кесектерінде: арқатөстік – 0,9 және 0,7 кг-ға, белдік – 0,5 және 0,3 кг-ға, жамбастық – 2,3 және 1,5 кг-ға ($P<0,05-0,001$). Ет ұшаларының органолептикалық, биохимиялық және спектрометрикалық көрсеткіштері бойынша өнімнің сапалылығы дәлелденді және де, сәйкесінше, сыналып отырған препараттардың қауіпсіздігі. Бұқалардың организмының биоресурстық әлеуетін іске асыру биопрепараттардың тұрақтылығына тән емес гемопоззаның, жасушалық және гуморальды факторлардың жандануы, ашық Prevention-N-A сәйкес эффектісін кезінде, орнатылды. Алынған мәліметтердің жаңалығы РФ өнертабыстарының Мемлекеттік реестрінде 26.10.2016 ж. және 19.06.2017 ж. сәйкес тіркелген өнертабыс патенттерімен № 2602687 және № 2622765 РФ расталған.

Түйін сөздер: бұқалар, өсіру, өсіріп жеткізу, бордақылау, Prevention-N-A және Prevention-N-E биопрепараттары, еттілік сапалары.

В. Г. Семенов¹, Д. А. Баймуханов², Н. И. Косяев¹, Р. М. Мударисов³,
Н. И. Морозова⁴, Ф. А. Мусаев⁴, Д. А. Никитин¹, М. Б. Калмагамбетов²

¹Чувашская государственная сельскохозяйственная академия, Чебоксары, Чувашская Республика, Россия,

²Казахский научно-исследовательский институт животноводства и кормопроизводства, Алматы, Казахстан,

³Башкирский государственный аграрный университет, Уфа, Республика Башкортостан, Россия,

⁴Рязанский государственный агротехнологический университет им. П. А. Костычева,
Рязань, Рязанская область, Россия

РОСТ, РАЗВИТИЕ И МЯСНЫЕ КАЧЕСТВА БЫЧКОВ НА ФОНЕ ПРИМЕНЕНИЯ БИОПРЕПАРАТОВ СЕРИИ PREVENTION

Аннотация. Впервые на основе комплексных исследований научно обоснована и экспериментально доказана зоотехническая целесообразность применения разработанных биопрепаратов Prevention-N-A и Prevention-N-E в технологии производства говядины для реализации биоресурсного потенциала мясных качеств бычков черно-пестрой породы. На фоне применения биопрепаратов установлена активизация роста и развития бычков в периоды выращивания, доращивания и откорма, что обусловило более высокие убойные и мясные качества туш и, как следствие, выход ценных отрубов: спиногорудного – на 6,1 и 4,0 кг ($P < 0,01-0,001$), поясничного – на 2,6 и 1,7 кг ($P < 0,05-0,01$) и тазобедренного – на 8,6 и 7,1 кг ($P < 0,001$), нежели контроле. Наибольшим содержанием мякоти высшего сорта характеризовались туши бычков 1-й (27,8±0,72 кг) и 2-й (26,7±0,58 кг) опытных групп соответственно на 3,5 и 2,4 кг по сравнению с контролем (24,3±0,73 кг), а также их отруба: спиногорудной – на 0,9 и 0,7 кг, поясничный – на 0,5 и 0,3 кг, тазобедренный – на 2,3 и 1,5 кг ($P < 0,05-0,001$). Доказана доброкачественность мясных туш по органолептическим, биохимическим и спектрометрическим показателям и, следовательно, безопасность испытуемых препаратов. Установлено, что реализация биоресурсного потенциала организма бычков была вызвана активизацией гемопоеза, клеточных и гуморальных факторов неспецифической устойчивости биопрепаратами, при более выраженном соответствующем эффекте Prevention-N-A. Новизна полученных данных подтверждена патентами РФ на изобретение № 2602687 и № 2622765, зарегистрированных в Государственном реестре изобретений РФ 26.10.2016 г. и 19.06.2017 г. соответственно.

Ключевые слова: бычки, выращивание, доращивание, откорм, биопрепараты Prevention-N-A и Prevention-N-E, мясные качества.

About the authors:

Seменов Vladimir Grigoryevich, doctor of biological science, professor, honored worker of science of the Chuvash Republic, professor of department of morphology, obstetrics and therapy of the Chuvash state agricultural academy, Cheboksary, Chuvash Republic, Russia, e-mail: semenov_v.g@list.ru.

Baimukanov Dastanbek Asylbekovich, doctor of agricultural sciences, professor, corresponding member of National Academy of Sciences of the Republic of Kazakhstan, chief researcher of department of cultivation and selection of the dairy cattle of the Kazakh research institute of livestock production and forage production, Almaty, Republic of Kazakhstan, e-mail: dbaimukanov@mail.ru.

Kosyaev Nikolay Ivanovich, doctor of veterinary sciences, professor, professor of department of an epizootology, parasitology and veterinary and sanitary examination of the Chuvash state agricultural academy, Cheboksary, Chuvash Republic, Russia, e-mail: kocyevni81@mail.ru.

Mударисов Rinat Mansafovich, doctor of agricultural sciences, professor, professor of department of beekeeping, private zootechnics and animal husbandry of the Bashkir state agricultural university, Ufa, Republic of Bashkortostan, Russia, e-mail: r-mударисов@mail.ru.

Morozova Nina Ivanovna, doctor of agricultural sciences, professor, honored worker of the higher school of the Russian Federation, head of the department of the production technology and processing of agricultural production of the Ryazan state agrotechnological university of P.A. Kostychev, Ryazan, Ryazan region, Russia, e-mail: morozova@rgatu.ru.

Mусаев Farrukh Ataullovich, doctor of agricultural sciences, professor, professor of department of the production technology and processing of agricultural production of the Ryazan state agrotechnological university of P.A. Kostychev, Ryazan, Ryazan region, Russia, e-mail: musaev@rgatu.ru.

Никитин Dmitry Anatolyevich, candidate of veterinary sciences, associate professor of morphology, obstetrics and therapy of the Chuvash state agricultural academy, Cheboksary, Chuvash Republic, Russia. e-mail: nikitin_d_a@mail.ru.

Калмагамбетов Murat Baitugelovich, Candidate of Agricultural Sciences, Head of the Department of Feeding Livestock Animals, Limited Liability Partnership “Kazakh Scientific Research Institute of Animal Breeding and Fodder Production”. E-mail: animal_feeding@mail.ru.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-1467 (Online), ISSN 1991-3494 (Print)

<http://www.bulletin-science.kz/index.php/ru/>

Редакторы *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 06.03.2018.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
14,2 п.л. Тираж 500. Заказ 2.