

ISSN 2518-1467 (Online),
ISSN 1991-3494 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Х А Б А Р Ш Ы С Ы

ВЕСТНИК

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

THE BULLETIN

THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

PUBLISHED SINCE 1944

3

MAY – JUNE 2019

ALMATY, NAS RK

NAS RK is pleased to announce that Bulletin of NAS RK scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of Bulletin of NAS RK in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential multidiscipline content to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабаршысы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабаршысының Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді мультидисциплинарлы контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Вестник НАН РК» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Вестника НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному мультидисциплинарному контенту для нашего сообщества.

Б а с р е д а к т о р ы

х. ғ. д., проф., ҚР ҰҒА академигі

М. Ж. Жұрынов

Р е д а к ц и я а л қ а с ы:

Абиев Р.Ш. проф. (Ресей)
Абишев М.Е. проф., корр.-мүшесі (Қазақстан)
Аврамов К.В. проф. (Украина)
Аппель Юрген проф. (Германия)
Баймуқанов Д.А. проф., корр.-мүшесі (Қазақстан)
Байтулин И.О. проф., академик (Қазақстан)
Банас Иозеф проф. (Польша)
Берсимбаев Р.И. проф., академик (Қазақстан)
Велесько С. проф. (Германия)
Велихов Е.П. проф., РҒА академигі (Ресей)
Гашимзаде Ф. проф., академик (Әзірбайжан)
Гончарук В.В. проф., академик (Украина)
Давлетов А.Е. проф., корр.-мүшесі (Қазақстан)
Джрбашян Р.Т. проф., академик (Армения)
Қалимолдаев М.Н. проф., академик (Қазақстан), бас ред. орынбасары
Лаверов Н.П. проф., академик РАН (Россия)
Лупашку Ф. проф., корр.-мүшесі (Молдова)
Мохд Хасан Селамат проф. (Малайзия)
Мырхалықов Ж.У. проф., академик (Қазақстан)
Новак Изабелла проф. (Польша)
Огарь Н.П. проф., корр.-мүшесі (Қазақстан)
Полещук О.Х. проф. (Ресей)
Поняев А.И. проф. (Ресей)
Сагиян А.С. проф., академик (Армения)
Сатубалдин С.С. проф., академик (Қазақстан)
Таткеева Г.Г. проф., корр.-мүшесі (Қазақстан)
Умбетаев И. проф., академик (Қазақстан)
Хрипунов Г.С. проф. (Украина)
Юлдашбаев Ю.А. проф., РҒА корр.-мүшесі (Ресей)
Якубова М.М. проф., академик (Тәжікстан)

«Қазақстан Республикасы Ұлттық ғылым академиясының Хабаршысы».

ISSN 2518-1467 (Online),

ISSN 1991-3494 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы»РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5551-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 2000 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://www.bulletin-science.kz/index.php/en/>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д. х. н., проф. академик НАН РК
М. Ж. Журинов

Р е д а к ц и о н н а я к о л л е г и я:

Абиев Р.Ш. проф. (Россия)
Абишев М.Е. проф., член-корр. (Казахстан)
Аврамов К.В. проф. (Украина)
Апель Юрген проф. (Германия)
Баймуканов Д.А. проф., чл.-корр. (Казахстан)
Байтулин И.О. проф., академик (Казахстан)
Банас Иозеф проф. (Польша)
Берсимбаев Р.И. проф., академик (Казахстан)
Велесько С. проф. (Германия)
Велихов Е.П. проф., академик РАН (Россия)
Гашимзаде Ф. проф., академик (Азербайджан)
Гончарук В.В. проф., академик (Украина)
Давлетов А.Е. проф., чл.-корр. (Казахстан)
Джрбашян Р.Т. проф., академик (Армения)
Калимолдаев М.Н. академик (Казахстан), зам. гл. ред.
Лаверов Н.П. проф., академик РАН (Россия)
Лунашку Ф. проф., чл.-корр. (Молдова)
Моход Хасан Селамат проф. (Малайзия)
Мырхалыков Ж.У. проф., академик (Казахстан)
Новак Изабелла проф. (Польша)
Огарь Н.П. проф., чл.-корр. (Казахстан)
Полещук О.Х. проф. (Россия)
Поняев А.И. проф. (Россия)
Сагиян А.С. проф., академик (Армения)
Сатубалдин С.С. проф., академик (Казахстан)
Таткеева Г.Г. проф., чл.-корр. (Казахстан)
Умбетаев И. проф., академик (Казахстан)
Хрипунов Г.С. проф. (Украина)
Юлдашбаев Ю.А. проф., член-корр. РАН (Россия)
Якубова М.М. проф., академик (Таджикистан)

«Вестник Национальной академии наук Республики Казахстан».

ISSN 2518-1467 (Online),

ISSN 1991-3494 (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5551-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год

Тираж: 2000 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18.

www: nauka-nanrk.kz, bulletin-science.kz

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

E d i t o r i n c h i e f

doctor of chemistry, professor, academician of NAS RK

M. Zh. Zhurinov

E d i t o r i a l b o a r d:

Abiyev R.Sh. prof. (Russia)
Abishev M.Ye. prof., corr. member. (Kazakhstan)
Avramov K.V. prof. (Ukraine)
Appel Jurgen, prof. (Germany)
Baimukanov D.A. prof., corr. member. (Kazakhstan)
Baitullin I.O. prof., academician (Kazakhstan)
Joseph Banas, prof. (Poland)
Bersimbayev R.I. prof., academician (Kazakhstan)
Velesco S., prof. (Germany)
Velikhov Ye.P. prof., academician of RAS (Russia)
Gashimzade F. prof., academician (Azerbaijan)
Goncharuk V.V. prof., academician (Ukraine)
Davletov A.Ye. prof., corr. member. (Kazakhstan)
Dzhrbashian R.T. prof., academician (Armenia)
Kalimoldayev M.N. prof., academician (Kazakhstan), deputy editor in chief
Laverov N.P. prof., academician of RAS (Russia)
Lupashku F. prof., corr. member. (Moldova)
Mohd Hassan Selamat, prof. (Malaysia)
Myrkhalykov Zh.U. prof., academician (Kazakhstan)
Nowak Isabella, prof. (Poland)
Ogar N.P. prof., corr. member. (Kazakhstan)
Poleshchuk O.Kh. prof. (Russia)
Ponyaev A.I. prof. (Russia)
Sagiyani A.S. prof., academician (Armenia)
Satubaldin S.S. prof., academician (Kazakhstan)
Tatkeyeva G.G. prof., corr. member. (Kazakhstan)
Umbetayev I. prof., academician (Kazakhstan)
Khripunov G.S. prof. (Ukraine)
Yuldashbayev Y.A., prof. corresponding member of RAS (Russia)
Yakubova M.M. prof., academician (Tadjikistan)

Bulletin of the National Academy of Sciences of the Republic of Kazakhstan.

ISSN 2518-1467 (Online),

ISSN 1991-3494 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5551-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 2000 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/>, <http://bulletin-science.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

**Yu. A. Yuldashbayev¹, M. I. Selionova², M. M. Aibazov², S. I. Svetlichny², N. N. Bondarenko¹,
S. V. Svistunov¹, D. A. Baimukanov⁴, S. O. Chylbak-ool¹, A. A. Tlepov⁵**

¹FSBEI HE Russian State Agrarian University – Moscow Agricultural Academy named after K. A. Timiryazev, Moscow, Russia,

²Kuban State Agrarian University, Krasnodar, Russia,

³All-Russian Scientific Research Institute of Sheep and Goat breeding – branch of the FSBSI North Caucasus Federal Scientific Agrarian Center, Stavropol, Russia,

⁴Kazakh National Agrarian University, Almaty, Kazakhstan,

⁵Taraz State University named after M. Kh. Dulati, Taraz, Kazakhstan.

E-mail: zoo@rgau-msha.ru, m_selin@mail.ru, m_selin@mail.ru, s.s.i.23@yandex.ru, bondarienko.49@mail.ru, svistunov@list.ru, dbaimukanov@mail.ru, zoo@rgau-msha.ru, tlepov_anarbek@mail.ru

ESTRUS INDUCTION IN DAIRY SHEEP DURING THE ANESTROUS PERIOD

Abstract. The aim of the research was to investigate the possibility of estrus induction in dairy sheep during the anestrus period using various hormonal preparations and finding out the effectiveness of this biotechnological method.

Experiments on the estrus induction were conducted in the Nikolaev M.I. peasant farm of Krymsky region, Krasnodar Krai. The object of the research was lactating and dry sheep of the dairy breed "Lacaune" with a total number of 332 heads between the ages of 2.5 and 5 years.

It is shown that in the anestrus season (from February to August) in the climatic conditions of the Krasnodar Krai with sufficiently high effectiveness (54.3-63.2%), it is possible to induce estrus in the Lacaune dairy sheep breed. In this case, the object of stimulating estrus can be both lactating and dry females. High fertility (75.0%) in sheep with induced estrus was established in the summer period. In the winter and spring seasons, this figure was 69.3% and 60.7%, respectively. The greatest effectiveness of estrus induction was established during hormonal stimulation of sheep during the summer period when the largest number of lambs (0.71) per one ewe was received. In the spring and winter seasons, this indicator was 0.37 and 0.58, respectively.

Effective technology of hormonal estrus induction in the unsexual season allows for the planned reproduction of dairy sheep throughout the year for year-round rhythmic milk production.

Key words: dairy ewes, anestrus period, estrus cycle, hormone therapy, estrus induction, progestagens, insemination, fertility.

Introduction. Sheep are polycyclic animals with a pronounced sexual season. Outstanding scientist A.I. Lopyrin believed that in the northern hemisphere, sheep start to be tugging in 8-10 weeks after the longest day of light. Therefore, the sexual (estrus) season in almost all sheep breeds occurs in autumn, as a rule, from September to December [1]. At this time, they are inseminated, while the period of mass young stock production falls on February-May. In economic conditions, when the most important product of sheep farming was high-quality wool, which was sheared once a year, the seasonality of sheep breeding was not decisive. Currently, in connection with the transfer of sheep farming to the priority meat direction and the initiation of dairy sheep farming in the country, the biologically determined seasonality of sheep breeding has become a critical point limiting the increase in the efficiency of the industry. This is due to the fact that the market requires the receipt of meat and dairy products and their delivery throughout the year [2, 3]. Accordingly, the technology for year-round sheep reproduction is needed.

Scientists and experts all over the world try to overcome the biological barrier to the seasonality of sheep breeding. As a rule, they are divided into two directions. The first is sheep breeding improvement in

order to ensure polyestrous, i.e. the ability to be tuppings and become fertile at any time of the year was a genetically determined trait. Such fairly common breeds as the Romanov, Il-de-France, Dorper, Finnish Landrace, East Frisian, Merinoland and, partly, Dorset have practically no breeding season. However, the process of breeding polyestrous sheep breeds is long and it is not always possible to get a predictable result. In addition, even in polyestrous sheep breeds, there is significant variability of cyclicity in the seasons of the year. [4].

The second direction is the use of various factors of external influence on the female's organism, based on a detailed knowledge of the mechanisms of the neurohumoral regulation of the reproductive function in sheep.

In order to induce estrus in sheep during the anestrus period, two methods are used - light and hormonal, each of which has its own advantages and disadvantages, determining the appropriateness and extent of their use in specific economic conditions.

The light method of estrus induction in sheep is based on the fact that the hypothalamic centers that regulate the release of gonadotropic hormones are adapted to light effects and are stimulated by the natural light difference when changing a long summer day to a shorter autumn one. Creating a so-called "artificial autumn", as shown by numerous experiments, stimulates estrus in some sheep [5]. However, the majority of authors agree that the number of sheep that have been tuppings varies widely (from 16 to 80%), excluding the predicted and successful application of this method with a practically significant result. [1].

The classic method of estrus induction in the off-season is the induction of estrus with the help of hormonal drugs, based on the prolongation of the luteal phase of the estrus by progestagens [1, 6]. Some authors mistakenly believe that "progesterone, affecting the pituitary gonadotropic cells, stimulates the synthesis and release of follicle-stimulating hormone (FSH) and luteinizing hormone (LH), which are involved in the regulation of ovarian function and the formation of follicles" [4]. On the contrary, progestagens inhibit the secretory function of the hypothalamus and the release of releasing hormone, which leads to suppression of the function of the posterior pituitary and release of follicle-stimulating hormone. Low concentration of FSH and LH in the peripheral blood leads to inhibition of the generative function of the ovaries, respectively, the growth and development of follicles are suppressed. As a result of hormonal treatment with progestagens, in animals, the stages of the estrus cycle are equalized, and after their action ceases, the physiological status of the generative apparatus of the females comes close to the post-luteal (or preovulatory) phase of the estrus cycle, the gonadotropic cells of the posterior lobe of hypophysis are activated, and the growth and maturation of the follicles in the ovaries are enhanced in all treated animals and, accordingly, animals are simultaneously tuppings [7-9].

As noted above, year-round rhythmic milk production is the main technological aspect of effective dairy sheep farming. This requires the breeding of dairy sheep in both the estrous and anestrus periods. However, for different breeds of sheep, their age and physiological state, the adopted technology of production in different climatic conditions, hormonal treatment to induce estrus and its effectiveness may vary significantly [8,9,10]. In this regard, the study of the possibility of estrus induction in dairy sheep and finding out the effectiveness of this biologically-inspired technique is relevant, which determined the aim of this investigation.

Materials and research methods. Own experiments on the estrus induction were conducted in the Nikolaev M.I. peasant farm of the Krymsky district of Krasnodar Krai (figure 1). The object of the research was lactating and dry sheep of the dairy breed "Lacaune". For the induction of estrus, several drugs were used, injected and intravaginally. For the treatment of one group of sheep, the common drug "progesterone" was used, which was administered intramuscularly at a dose of 0.7 ml for 11 days. For hormonal treatment of two other groups of sheep, intravaginal pessaries of domestic production were used, impregnated with AMOL, the active ingredient of which is the synthetic analogue of progesterone 17 α -mepregnenol acetate and imported Syncro-part pessaries (France), containing 30 mg of active ingredient flugestone acetate (Flugestone acetatum).

In all groups of sheep, to stimulate the growth of follicles in the ovaries after the removal of the progestogenic preparation, pregnant-mare serum (PMS) (Syncro-part Pmsg 6000me France) and the Follimag drug were used in different doses. Studies were conducted from February to August. In order to determine the impact of the season on the effectiveness of estrus induction, experiments with a certain degree of conditionality were divided into winter (February-March), spring (April-May) and summer (June-August).



Figure 1 – Interior design of the premises for the maintenance and milking of the Lacaune breed sheep in Nikolaev M.I. peasant farm of Krasnodar Krai

The methodological features of individual experiments are given in the description of experimental results.

Results and discussion.

Experiment 1. The first experiment was conducted in February-March 2017. 3 groups of animals were formed.

In the first group of sheep (lactating ewes in the amount of 85 heads), pessaries impregnated with the AMOL preparation at a dose of 30 mg AS were intravaginally administered for 12 days. After sponge removal, all the animals were intramuscularly injected with PMS at a dose of 600 units.

The second group of sheep was formed from dry ewes in the number of 51 heads, which were treated similarly to group 1.

For the treatment of the third group sheep, also consisting of dry females ($n = 17$), the “progesterone” drug was used, which was administered intramuscularly at a dose of 0.7 ml for 11 days. After treatment with progestagens in order to stimulate folliculogenesis, the Follimag drug was administered once intramuscularly at a dose of 500 units.

In all groups of sheep, sampling in estrus was not carried out, insemination of sheep was conducted after 55 hours, using freshly obtained semen for quality that meets the minimum requirements of the “Manual on the operating procedures of organizations of artificial insemination and transplantation of embryos of farm animals” (M., 2000).

The following results were attained. In the first group, 61 sheep were fertilized (71.2%), 62 lambs were born (fertility - 101.6%). In the second group, 34 sheep (66.6%) were fertilized, from which 36 lambs were obtained (fertility - 105.8%). In the third group, insemination performance was 64.7% (11 out of 17 inseminated sheep were fertilized), from which 11 lambs were obtained (fertility 100%).

Thus, in winter, out of 153 treated females, 106 sheep were fertilized fruitfully (average fertility 69.3%), from which 109 lambs were obtained (average fertility 102.8%). There was no significant difference in fertility between lactating and dry ewes. The average number of lambs obtained per 1 treated sheep was 0.71.

Experiment 2. The second experiment was performed in April - May 2017. Two groups of sheep were formed.

The first group, consisting of lactating ewes ($n = 58$), was treated with the intravaginal injection of sponges impregnated with the AMOL preparation at a dose of 30 mg AS. The intravaginal exposure period of progestagens was 12 days.

The second group ewes (dry ewe, $n = 45$) were also intravaginally injected for 12 days with Syncro-part pessaries (France) containing 30 mg of active substance of fluogestone acetate (Flugestone acetatum).

After removing the sponges, all animals were intramuscularly injected one time: in the first group, PMS was administered at a dose of 600 units. (Syncro-part Pmsg 6000me France), in the second group - “Follimag” at a dose of 500 units.

In order to carefully establish the estrus-inducing effect of the drugs used in this experiment, after the administration of gonadotropin, sheep were controlled for the state of estrus. To do this, with the help of sheep-samplers with tied up aprons, sheep were selected in the estrus. For a more accurate determination of the estrus onset, the sample was carried out every 4 hours.

The following results were obtained.

In the first group of 58 sheep, during the first 20 hours after the PMS injection, no one ewe has been tugging. After sampling after 24 hours, 2 heads in estrus were selected, after 28 hours - 3 heads, after 32 hours - 6 heads, after 36 hours - 10 heads, after 40 hours - 11 heads, after 44 hours - 2 heads. Sampling after 48 hours and further till insemination, which was carried out 55-56 hours after the PMS administration, did not give positive results.

Therefore, in the first group of 58 lactating ewes, after progestagens + PMS treatment, 34 sheep were tugging within 44 hours (58.6%). At the same time, of these 34 animals, 27 sheep (79.4%) have been tugging in a period from 28 to 40 hours after the gonadotropin administration.

All animals in the estrus were inseminated with freshly obtained semen with a mobility of 7-8 points, a concentration of 2.6-2.8 billion/ml. Of the inseminated 34 heads, 22 sheep lambed (fertilization rate was 64.7%). 24 lambs were born (fertility was 109.1%).

In such a way, analyzing the results obtained in the first group of animals, we came to the conclusion that, out of 58 lactating ewes, after the estrus induction in spring, 22 heads were fruitfully inseminated (37.9%), with a yield of 0.41 lamb per 1 treated ewe.

In the second group of 45 sheep, within the first 24 hours after the PMS injection, no one single ewe was tugging. After sampling, after 28 hours, 3 blissom sheep were selected, after 32 hours - 3 heads, after 36 hours - 13 heads, after 40 hours - 2 heads, after 44 hours - 1 head. As in the first group, 48 hours later, before insemination, which was carried out 55-56 hours after the Follimag administration, the sample did not reveal blissom ewes.

In this group of 45 dry ewes, after treatment with progestogen + Follimag, 22 heads were tugging within 44 hours. (48.9%). At the same time, out of these selected 22 sheep in estrus, 21 animals (95.5%) have been tugging between 28 and 40 hours after administration of gonadotropin.

As in the first group, all the sheep in the estrus were inseminated with freshly obtained semen with a mobility of 8-9 points, a concentration of 2.8-3.0 billion/ml. Among 22 animals, 12 sheep lambed (fertilization rate was 54.5%). 14 lambs were born (fertility was 116.6%).

In such a manner, in the second group of animals of 45 dry ewes, after the estrus induction in the spring period, 12 heads were fruitfully inseminated (26.6%), with a yield of 0.31 lamb per 1 treated ewe.

For a more objective assessment of the effectiveness of estrus induction in sheep in spring, we found it possible to summarize the data obtained for two groups and to derive the averages. Of 103 estrus-induced ewes, estrus was recorded in 56 animals (54.3%), of which, after insemination, 34 sheep lambed (fertility rate was 60.7%). 38 lambs were obtained (fertility was 111.8% for a sheep that gave birth, or 65.5% for an inseminated sheep). The estrus induction in the spring season resulted in 0.37 lamb per treated sheep.

Experiment 3. The third experiment was conducted in June-August 2017. As in the second experiment, 2 groups of sheep were formed.

The first group, consisting of lactating ewes (n=36), was treated with the intravaginal injection of sponges impregnated with the "AMOL" at a dose of 30 mg AS. The progestagens exposure period was 14 days.

The second group of ewes (dry ewe, n = 40) was also intravaginally injected for 14 days with pessaries Syncro-part (France) containing 30 mg of fluogestone acetate (Flugestone acetatum).

After sponge removal, all animals were injected intramuscularly with gonadotropin: in the first group, PMS at a dose of 600 units. (Syncro-part Pmsg 6000me France), in the second group - "Follimag" at a dose of 400 units.

In the third experiment, as in the second one, in order to accurately determine the stimulating effect of the used drugs after the gonadotropin administration, sheep were sampled into the estrus state with the help of sheep samplers with tied up aprons. To accurately determine the onset of estrus, sampling was performed every 4 hours.

In this experiment, the following results were obtained.

In the first group of sheep, during the first 20 hours after the PMS injection, no one single ewe was tugging. After sampling after 24 hours, 1 head in estrus was selected, after 28 hours - 2 heads, after 32 hours - 7 heads, after 36 hours - 6 heads, after 40 hours - 5 heads after 44 hours and then before insemination, which was carried out 55-56 hours after the PMS administration, ewes in estrus were not detected.

Thus, in the first group of 36 sheep, after estrus induction, within 40 hours, 21 heads were tugging (58.3%). At the same time of these 21 heads, in the time period from 28 to 40 hours, estrus was induced in 20 sheep (95.2%) after the PMS administration. All animals in estrus were inseminated with freshly obtained semen with a mobility of 8-8.5 points, a concentration of 2.5-3.0 billion/ml. Among 21 sheep, 15 animals lambled (fertility rate was 71.4%). 18 lambs were obtained (fertility was 120.0%). Consequently, in the first group of animals of 36 lactating ewes, after the induction of estrus in the summer season, 15 of them were fruitfully inseminated (41.6%), in this case, 0.5 lamb was obtained per 1 treated ewe.

In the second group, sheep in estrus were detected only 28 hours after the Follimag administration, when 2 heads in a state of estrus were selected. After 32 hours, 2 heads were selected, after 36 hours - 10 heads, after 40 hours - 12 heads, after 44 - 1 head. Sampling after 48 hours and further did not reveal any ewes in estrus. In this group of 40 dry ewes, after progestagens + Follimag treatment, 27 animals were tugging within 44 hours (67.5%). At the same time, 26 selected sheep in estrus (96.2%) were tugging in the time period from 28 to 44 hours after the Follimag use.

As in the first group, all the sheep in estrus were inseminated with freshly obtained semen, the quality of which corresponded to the minimum requirements of the Instruction. Of 27 sheep, 21 animals lambled (fertilization rate was 77.7%). 26 lambs were born (fertility was 123.8%).

Thus, in the second group of animals of 40 dry ewes, after the estrus induction in the summer, 21 animals were fruitfully inseminated (52.5%), 0.65 lamb was obtained per 1 treated ewe.

As in the second experiment, we think it possible to summarize the data obtained in two groups and to derive the averaged indicators of the effectiveness of estrus induction in sheep in the summer period. Of 76 estrus-induced ewes, estrus was detected in 48 heads (63.2%), of which, after insemination, 36 sheep lambled (fertilization rate - 75.0%). A total of 44 lambs were obtained (fertility - 122.2% in terms of lambled sheep, or 0.9 lamb per inseminated ewe) (figure 2). The induction of estrus in the summer season resulted in a yield of 0.58 lamb per treated sheep.

The summarized data for the three experiments are presented in table.



Figure 2 – Breeding, estimation of growth and development of lambs got from ewes with induced estrus in the summer period

Indicators of the effectiveness of estrus induction in "Lacaune" dairy sheep breed in the anestrus period by the seasons

Season	Treated sheep, heads	were tugging		Of them lambled		Number of lambs		Obtained lambs per 1 treated sheep
		heads	%	heads	%	heads	%	
Winter (February-March)	153	—*	—*	106	69.3	109	102.8	0.71
Spring (April-May)	103	54	54.3	34	60.7	38	111.8	0.37
Summer (June-August)	76	48	63.2	36	75.0	44	122.2	0.58

*Sampling of ewes in estrus was not carried out. Frontal insemination of all sheep was used 55 hours after finishing the treatment with progestagens.

Comparison of data shows that the greatest fertilization rate (75.0%) in sheep with induced estrus was established in the summer period. In the winter and spring seasons, this figure was 69.3% and 60.7%, respectively.

Conclusions. The obtained results indicate that in the anestrus season (from February to August) in the climatic conditions of the Krasnodar Krai, with high efficiency, it is possible to induce estrus in the sheep of the Lacaune dairy breed. In this case, the object of estrus induction can be both lactating and dry females. At the same time, some of the resulted indicators, in our opinion, require more careful analysis in order to make possible adjustments in terms and estrus induction scheme in sheep.

The best fertilization rate of sheep during induced estrus was found in the summer period. The smaller value of this indicator was recorded in the winter period and the smallest - in the spring. However, it should be noted that in the winter experiment, no sheep in estrus were sampled, the frontal insemination was applied 55 hours after the administration of the gonadotropic hormone. In this regard, it is difficult to judge the effectiveness of estrus induction. At the same time, the main resulting indicator of our experiments in the winter period is 0.71 lamb per treated sheep. It can be assumed that a sufficiently high rate of fruitful insemination is connected with the fact that the winter group of animals could get a part of the sheep that were before the very end of the estrous season, in which, apparently, the natural hormonal background allowed to sensitize the sexual centers with endogenous progesterone, and the administration of exogenous progesterone (vaginal pessaries) coincided with the preovulatory phase of the estrus cycle, which led to the manifestation of estrus and successful fertilization.

In a similar way, it is possible to explain the high fertilization rate of females (69.3%) in the summer experiment (June-August). It could also happen that sheep with regenerating natural hormonal background (especially in August) fell into this group, which, after saturation of the body with exogenous progesterone followed by stimulation of folliculogenesis in the ovaries, showed a full estrus which was confirmed by obtaining 0.58 lamb per one treated sheep.

The smallest number of lambs (0.37 per one treated ewe) was obtained in the spring (April-May) treatment period. Apparently, out of the estrous season, the function of the pituitary and ovaries under the influence of unfavorable environmental factors (prolonged light day, insolation) noticeably weakens and even under the influence of exogenous gonadotropic hormone (PMS or Follimag), the developing ovarian follicles do not reach ovulation maturity and undergo partial luteinization without rupture and release of the egg. It may also happen that during this period so-called abortive ovulation when the selected eggs are defective and cannot be fertilized.

It is believed that in lactating females, the effectiveness of estrus induction and fertilization rate is lower than in dry (non-lactating) animals [6]. In our experiments, this pattern is quite clearly seen only during hormonal treatment in the summer.

Obviously, the obtained data are preliminary, and their analysis is not exhaustive. Moreover, the reproductive indicators obtained in our experiments are significantly lower than in the experiments of other authors [6]. In part, this can be explained by the fact that the data on fertility in these authors are given not by lambing, but by the early diagnosis of pregnancy of ewe by the ultrasound method, i.e. without taking into account possible fetal mortality. It is also possible the impact that the experiments were conducted in other climatic conditions and with other sheep breeds using various induction schemes [11-15]. For a more complete analysis and justification of the laws obtained, further research is needed, first of all, of the natural hormonal background in sheep of different age groups and physiological state in the context of the seasons of the year. Selective laparoscopy (using endoscopic equipment) to assess the state of the internal genital organs in treated animals could be a good diagnostic test to determine the effectiveness of estrus induction.

Ю. А. Юлдашбаев¹, М. И. Селионова², М. М. Айбазов², С. И. Светличный²,
Н. Н. Бондаренко¹, С. В. Свистунов¹, Д. А. Баймұқанов⁴, С. О. Чылбак-оол¹, А. Тлепов⁵

¹Жоғары білім беру саласындағы федералдық мемлекеттік бюджеттік білім беру саласының мемлекеттік орталығы – К. А. Тимирязев атындағы Мәскеу аграрлық академиясы, Мәскеу, Ресей,

²Кубань мемлекеттік аграрлық университеті, Краснодар, Ресей, ³Бүкілресейлік қой және ешкінің ғылыми-зерттеу институты – «Солтүстік-Кавказ федералдық ғылыми зерттеу орталығы» федералды мемлекеттік бюджеттік ғылыми институты филиалы, Ставрополь, Ресей,

⁴Қазақ ұлттық аграрлық университеті, Алматы, Қазақстан,

⁵М. Х. Дулати атындағы Тараз ұлттық университеті, Тараз, Қазақстан

АНЭСТРАЛЬДЫ КЕЗЕНДЕ СҮТТІ ҚОЙЛАРДЫҢ ЭСТРУС ИНДУКЦИЯСЫ

Аннотация. Зерттеудің мақсаты әртүрлі гормоналды препараттарды қолдану арқылы анэстральды кезеңде сүт қойларында сүтқоректілерді индукциялау мүмкіндігін зерттеу және биотехнологиялық әдістердің тиімділігін анықтау болды.

Жыныстық күйіт ынталандыру жөніндегі эксперименттер Қырым аймағы Краснодар өлкесінің Николаев М.И. Шаруа (фермер) шаруашылығында өтті. Зерттеудің нысаны 2,5 жылдан 5 жылға дейінгі кезеңде жалпы 332 басы бар сүт тұқымының «Лакон» ("Lacaune") лактация кезеңінде және суалу кезеңіндегі қойлары болды.

Краснодар өлкесінің климаттық жағдайында анэстральдық маусымда (ақпаннан тамызға дейін) жеткілікті жоғары тиімділікпен (54,3-63,2%) сүт қойларында «Лакон» күйіт табуға болады деп көрсетілген. Бұл жағдайда эструсты ынталандыратын зат сүтті және суалу кезеңіндегі аналық болуы мүмкін. Жазғы кезеңде индуцирленген күйіт кезде қойлардағы жоғары өнімділік (75,0%) белгіленді. Қысқы және көктемгі мезгілдерде бұл көрсеткіш тиісінше 69,3% және 60,7% құрады. Күйітті ынталандырудың ең тиімділігі жазғы кезеңде қойлардың гормоналды ынталандыруы кезінде анықталды, ол бір қойға ең көп қозы (0,71) алынды. Көктемгі-қысқы мезгілдерде бұл көрсеткіш тиісінше 0,37 және 0,58 құрады.

Мерзімді кезеңде эструстың гормоналды индукциясының тиімді технологиясы жыл бойына ритмді сүтті өндіруге сүт қойларын жоспарлы көбейтуге мүмкіндік береді.

Түйін сөздер: сүт аналықтары, анэстральдық кезеңі, эстрогендік цикл, гормондық терапия, эструс индукциясы, прогестогендер, ұрықтандыру, өнімділік.

Ю. А. Юлдашбаев¹, М. И. Селионова², М. М. Айбазов², С. И. Светличный²,
Н. Н. Бондаренко¹, С. В. Свистунов¹, Д. А. Баймуқанов⁴, С. О. Чылбак-оол¹, А. Тлепов⁵

¹ФГБОУ ВО РГАУ – МСХА им. К. А.Тимирязева, Москва, Россия,

²Кубанский государственный аграрный университет, Краснодар, Россия,

³Всероссийского НИИ овцеводства и козоводства – филиала ФГБНУ «Северо-Кавказский ФНАЦ», Ставрополь, Россия,

⁴Казахский национальный аграрный университет, Алматы, Казахстан,

⁵Таразский государственный университет им. М. Х. Дулати, Тараз, Казахстан

ИНДУКЦИЯ ЭСТРУСА У МОЛОЧНЫХ ОВЕЦ В АНЭСТРАЛЬНЫЙ ПЕРИОД

Аннотация. Целью исследования являлось изучение возможности индукции эструса у молочных овец в анэстральный период применением различных гормональных препаратов и выяснение эффективности этого биотехнологического приема.

Эксперименты по стимуляции половой охоты проведены в КФХ Николаев М.И. Крымского района Краснодарского края. Объектом исследований были лактирующие и сухостойные овцы молочной породы «Лакон» ("Lacaune") общей численностью 332 гол. в возрасте от 2,5 до 5 лет.

Показано, что в анэстральный сезон (с февраля по август) в природно-климатических условиях Краснодарского края с достаточно высокой эффективностью (54,3-63,2%) можно индуцировать половую охоту у овец молочной породы «Лакон». При этом объектом стимулирования эструса могут быть как лактирующие, так и сухостойные самки. Высокая оплодотворяемость (75,0%) у овец с индуцированной охотой установлена в летний период. В зимний и весенний сезоны этот показатель составил соответственно 69,3% и 60,7%. Наибольшая эффективность вызывания половой охоты установлена при гормональной стимуляции овец в летний период, когда было получено наибольшее число ягнят (0,71) на одну обработанную овцематку. В весенний и зимний сезоны этот показатель составил соответственно 0,37 и 0,58.

Эффективная технология гормонального индуцирования эструса в неполовой сезон позволяет проводить плановое размножение молочных овец в течение всего года для круглогодичного ритмичного получения молока.

Ключевые слова: молочные овцематки, анэстральный период, эстральный цикл, гормональная терапия, индукция эструса, прогестогены, осеменение, плодовитость.

Information about authors:

Yuldashbayev Yusupzhan Artykovich, Corresponding Member of the Russian Academy of Sciences, Doctor of Agricultural Sciences, Professor, Dean of the Faculty of Animal Science and Biology, Professor of the Department of Private Animal Science, Russian State University - Moscow Agricultural Academy named after K. A. Timiryazev, Moscow, Russia; zoo@rgau-msha.ru; <https://orcid.org/0000-0002-7150-1131>

Selionova Marina Ivanovna, Doctor of Biological Sciences, Professor of the RAS, Director of the All-Russian Scientific Research Institute of Sheep and Goat breeding – branch of the FSBSI North Caucasus Federal Scientific Agrarian Center, Stavropol, Russia; m_selin@mail.ru

Aibazov Ali-Magomet Mussaevich, doctor of agricultural sciences, professor, head of the sheep farming department of the All-Russian Scientific Research Institute of Sheep and Goat breeding – branch of the FSBSI North Caucasus Federal Scientific Agrarian Center, Stavropol, Russia; m_selin@mail.ru

Svetlichnyy Sergey Ivanovich, degree-seeking student of the Kuban State Agrarian University, Chief Technologist of Agro Holding OOO, Krasnodar, Russia; s.s.i.23@yandex.ru; <https://orcid.org/0000-0003-1822-4080>

Bondarenko Nina Nikolaevna, Doctor of Agricultural Sciences, Professor at the Department of Parasitology, Veterinary-sanitary expert examination and Zoohygiene of the Kuban State Agrarian University, Krasnodar, Russia; bondarienko.49@mail.ru; <https://orcid.org/0000-0003-0232-4861>

Svistunov Sergey Vladimirovich, Candidate of Agricultural Sciences, Senior Researcher of the Department of Breeding and Genetics of Agricultural Animals, Krasnodar Scientific Center for Zootechnology and Veterinary Medicine, Krasnodar, Russia; svistunov@list.ru; <https://orcid.org/0000-0002-9098-9953>

Baimukanov Dastanbek Asylbekovich, Corresponding Member of the National Academy of Sciences of the Republic of Kazakhstan, Doctor of Agricultural Sciences, Professor of the Department of Physiology, Morphology, and Biochemistry named after academician N. U. Bazanova, NJSC “Kazakh National Agrarian University”, Almaty, Kazakhstan; dbaimukanov@mail.ru; <https://orcid.org/0000-0002-4684-7114>

Chylbak-ool Salbak Olegovna, post-graduate student of the Department of Private Animal Science, Russian State University - Moscow Agricultural Academy named after K. A. Timiryazev, Moscow, Russian Federation; zoo@rgau-msha.ru; <https://orcid.org/0000-0003-3799-9009>

Tleпов Anarbek Amreevich, candidate of Agricultural Sciences, Associate Professor of the Department of Biologi, M. Kh. Dulati Taraz State University, Taraz, Kazakhstan; tleпов_anarbek@mail.ru; <https://orcid.org/0000-0003-2947-3840>

REFERENCES

- [1] Lopyrin A.I. (1971) Sheep breeding biology. Moscow. 320 p. (in Russ.).
- [2] Selionov M.I., Bobryshova G.T., Gadzhiev Z.K., Izmailkov S.A. (2017) Sheep farming economy: pros and cons // Sheep, goats, wool business. N 1. P. 5-9 (in Russ.).
- [3] Aibazov M.M., Malakhova L.S., Trubnikova P.V. (2006) The results of the estrus induction in dairy goats in the anestrus period // Sheep, goats, wool business. N 2. P. 34-35 (in Russ.).
- [4] Tikhona G.S., Bezvesilnaya A.V., Khmelkov V.N. (2013) Influence of hormonal drugs on folliculogenesis in sheep in the anestrus period // Scientific and Technical Bulletin IT NAAN. N 109. P. 277-282 (in Russ.).
- [5] Kazakov V.M. (1962) On the artificial induction of estrous in sheep. Livestock, 1962. N 8 (in Russ.).
- [6] Chekunkova Yu.A. (2016) Stimulation of hunting in sheep in the spring period // Bulletin of the Altai State Agrarian University. N 8 (142) (in Russ.).
- [7] Karynbaev A.K., Akynbekova R. (2013) Effect of gonadotropic hormones on the production of eggs of the Karakul ewes of different ages // Sheep, goats, wool business. N 3. P. 31-32 (in Russ.).
- [8] Baimukanov D.A., Abugaliyev S.K., Seidaliyev N.B., Semenov V.G., Chindaliyev A.E., Dalibayev E.K., Zhamalov B.S., Muka Sh.B. (2019) Productivity and estimated breeding value of the dairy cattle gene pool in the Republic of Kazakhstan // Bulletin of National academy of sciences of the Republic of Kazakhstan. Vol. 1, N 377. P. 39-53. <https://doi.org/10.32014/2019.2518-1467.5> ISSN 2518-1467 (Online), ISSN 1991-3494 (Print).
- [9] Bogdan L., Groza I., Pop R., Petrean A., Bogdan S. (2011) Induction and oestrus synchronization in sheep during the breeding and nonbreeding season // Bulletin UASVM. Veterinary Medicine. 68 (2): 1843-5270.
- [10] Maraček I., Vlčková R., Kařatová J., Sopková D., Klapáčová K., Valocký I., Pořivák J. (2009) Effect of assisted oestrus on the ovulation rate and reproductive performance of Tsigai sheep // Slovak J. Anim. Sci. 2009. 42, SUPPLEMENT 1: 51-55 CVŽV Nitra ISSN 1337-9984.
- [11] Omontese B.O., Rekwot P.I., Rwuuan J.S., Ate I.U., H. Makun J. (2014) Induction of oestrus in Nigerian Ouda ewes with different oestrus synchrony protocols // Revue Méd. Vét. 165. 7-8, 240-244.
- [12] Stoycheva I., Kirilov A. (2015) Induction of synchronic oestrus, impregnancy and fertility of female lambs at 18 months of age and female lambs at 7-8 months of age, after treatment with PMSG // Bulgarian Journal of Agricultural Science. 21 (No 5). P. 1044-1048.
- [13] Hernández-Marín J.A., Cortez-Romero C., Corredor C.A.H. Sánchez J.G. (2018) Male effect and temporary weaning in synchronization of post-partum ovarian activity in Pelibuey ewes // South African Journal Of Animal Science. 48(4): 743-750. doi:10.4314/sajas.v48i4.16
- [14] Letelier C., Contreras I., Garcia-Fernandez R.A., Matildo Sanchez A.M. (2011) Effects of oestrus induction with progestagens or prostaglandin analogues on ovarian and pituitary function in sheep // Animal reproduction science. 126 (1-2): 61-9. Doi:10.1016/j.anireprosci.2011.04.012
- [15] Ashour G., El-Bassiony M.F., Dessouki Sh.M., El-Wakeel M.A. (2018) Application of different hormonal protocols for improving reproductive performance of Barki ewes // World's Veterinary Journal. 8 (3): 55-64.

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

ISSN 2518-1467 (Online), ISSN 1991-3494 (Print)

<http://www.bulletin-science.kz/index.php/en/>

Редакторы *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 10.06.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
12,7 п.л. Тираж 500. Заказ 3.